

**Deformation and metasomatism in the sub-continental lithospheric mantle  
of the Carpathian-Pannonian region (Hungary) and  
Jeju Island (South Korea)**

**Károly Hidas**

Summary of the Ph.D. dissertation

Ph.D. program for Geology and Geophysics at the Ph.D. school of Earth Sciences

Eötvös Loránd University, Budapest

Chair of the Ph.D. school: Prof. Miklós Monostori  
*(Department of Paleontology, Budapest)*

Advisor: Csaba Szabó, Ph.D. associate professor  
*(Department of Petrology and Geochemistry, Budapest)*

Tutors: György Falus, Ph.D.  
*(Eötvös Loránd Geophysical Institute of Hungary, Budapest)*  
István Kovács, Ph.D.  
*(Eötvös Loránd Geophysical Institute of Hungary, Budapest)*  
Prof. Yang Kyoung-hee  
*(Pusan National University, Busan, South Korea)*

2009 Budapest, Hungary

Keywords: fluid and silicate melt inclusion – metasomatism – deformation – rheology – upper  
mantle peridotite – Jeju Island, South Korea – Bakony-Balaton Highland Volcanic Field,  
Hungary

© 2009 – Hidas Károly  
All rights reserved.

## **Table of contents**

Introduction .....	2
Methodology .....	3
Results .....	5
Acknowledgements .....	7
Published papers and selected abstracts written in the framework of this Ph.D.....	7
References .....	10

## Introduction

The basic principle of plate tectonics and its relationship to volcanism is common knowledge yet in order to better understand in details one must consider the upper mantle as the origin and driving mechanism of these processes. Results of the past decades on melt generation, percolation and related metasomatic processes, along with the recognition that fabric analysis and crystallographic preferred orientation of constituent minerals has come to the foreground in recent studies on deep lithospheric rock samples inspired us to gain additional information on such well-known and widely inferred processes as deformation and metasomatism of the upper mantle, as well as melt-volatile immiscibility at high PT conditions. For such purposes, two peridotite xenolith series became available from the sub-continental lithospheric mantle of the Carpathian-Pannonian region and that of the Jeju Island showing clear evidences for either metasomatism with melt percolation or deformation, or both. The Carpathian-Pannonian region (CPR), which consists of a group of young extensional basins, in the eastern neighborhood of the Alpine orogen belt is situated on an extremely thinned continental lithosphere and it is one of the geologically best studied areas in the world (*e.g.* Csontos & Vörös, 2004, Horváth *et al.*, 2006, Bada *et al.*, 2007 and references therein). The fact that orogenic and extensional processes here are relatively young and that there is a massive body of knowledge on geology, geochemistry, geophysics, volcanism and sediments provides us a unique natural laboratory to study the evolution of the upper mantle. Besides, upper mantle peridotites from the sub-continental lithospheric mantle of the CPR are extensively studied (see Szabó *et al.*, 2004 for a summary), due to the Neogene alkali basaltic volcanism transporting large amount of xenoliths to the surface. Majority of these papers focuses on the central part of the CPR, particularly on the Bakony-Balaton Highland Volcanic Field (BBHVF), thus a broad geochemical and petrophysical database, as well as a solid knowledge has become available to outline the general composition and evolution of the sub-continental lithospheric mantle beneath the CPR. Nevertheless, special samples are needed to study deformation and metasomatism, two processes which widely occur in upper mantle and might significantly change its physico-chemical properties. We have selected two amphibole-bearing metasomatized spinel peridotites, which enclose coexisting silicate melt and fluid inclusions from the BBHVF, because they uniquely encapsulate metasomatism as is. Similarly, three deformed, flattened tabular equigranular spinel peridotites have been studied in details to better understand the deformation processes in the shallow sub-continental lithospheric mantle of the BBHVF. Although, the number and uniqueness of these selected xenoliths does not allow us to draw general conclusions on the physical state and geochemical composition of the sub-continental lithospheric mantle beneath the CPR, but comparing them to the general information already available might provide additional knowledge on the composition and evolution of the upper mantle. In contrast, Jeju Island is known as an intraplate volcano showing geochemical signatures of oceanic island basalt, whereas the island is located close to the eastern margin of the Eurasian plate which is known as a convergent plate margin, where Pacific and

Philippine plates are being subducted beneath the Eurasian Plate. Despite the fascinating geodynamic position of the island, peridotite studies from South Korea, particularly Jeju Island, are very rare (*e.g.* Choi *et al.*, 2002 and references therein). Hence, the general view on the evolution of the sub-continental lithospheric mantle here is still enigmatic. For this thesis we have selected a set of twenty peridotites, both common and special, to outline the general petrophysical feature of the shallow lithospheric mantle beneath the Island and to reveal special deformation and metasomatic processes which contributed to the composition and evolution of the upper mantle in East Asia.

This thesis is composed of the following major parts. (1) Because previous, especially, the early papers from the CPR described mainly the basic textural and geochemical features and solely presented “representative” xenoliths, only slight chemical and/or textural differences have been recognized. In this manuscript, we sought to provide additional information on the special rheological and textural and metasomatic processes and evolution of the lithospheric mantle beneath the central CPR by a detailed study on selected peridotite samples from a series of hundreds of xenoliths. By the reconstruction of pre- and post-entrapment evolution of silicate melt inclusions and fluid inclusions revealed from metasomatized peridotites of the CPR this study might contribute to the general knowledge of processes associated with high PT immiscibility and fluid entrapment. These peridotite xenoliths provide an opportunity to study the volatile content of such silicate melts, as well as the distribution of trace elements between melt and fluid during immiscibility in the upper mantle. These samples also show evidence regarding the presence of H<sub>2</sub>O in the silicate melt and fluid inclusions, in addition to the more common and abundant CO<sub>2</sub>. (2) Conversely, a reliable geochemical and textural database from peridotite xenoliths of Jeju Island, from which the general consequences on the evolution of the sub-continental lithospheric mantle can be drawn, is not yet available. Here, we present the first petrophysical analysis of “representative” and special mantle peridotites from the region along with their detailed major and trace element analysis to introduce deformation accompanied by geochemical evolution. (3) Comparison of the two peridotite series is necessary to reveal general features of the sub-continental lithospheric mantle and to distinguish them from those chemical and petrological features that are the result of the different geodynamic setting.

## **Methodology**

3 deformed and 2 metasomatized peridotite xenoliths were selected for detailed study from the alkali basaltic outcrops of the Bakony-Balaton Highland Volcanic Field (central Carpathian Pannonian region), near the villages of Szigliget and Szentbékállá. From Jeju Island (South Korea) 3 fine-grained mylonitic, 2 fine-grained porphyroclastic and 15 coarse grained protogranular-porphyroclastic peridotite xenoliths have been studied from three alkali basaltic localities at Sinsanri, Sangumburi and Jigriorem. 100-150  $\mu\text{m}$  thick doubly polished thin sections and separated doubly polished mineral grains were made for the different analytical purposes.

Olivine crystallographic preferred orientation (CPO) from the deformed peridotites of the BBHVF and 7 selected peridotites from Jeju Island was measured using a JEOL JSM-5600 scanning electron microscope at Geosciences Montpellier of Montpellier University 2 (Montpellier, France). This SEM is equipped with an electron back-scattered diffraction (EBSD) system from HKL Technology using Channel5 software. Accelerating voltage of 17 kV, spot size of 78 and 25 nm working distance were used. Automatic orientation mapping was carried out in at least a 10×10 mm area of the studied xenoliths. Depending on grain size, a step size between 12-100 microns was used. Data were evaluated using Channel 5 software package.

Major element composition of constituent minerals in the two peridotite series and the residual glass of silicate melt inclusions in the metasomatized peridotites of the BBHVF was measured by different electron microprobes (JEOL Superprobe JXA-8600, JEOL Superprobe JX-8200, Cameca SX-50, Cameca SX-100) equipped with WDS detectors at the Department of Earth Sciences, University of Florence (Italy); Department of Geosciences, Virginia Polytechnic Institute and State University (USA); Bayerisches Geoinstitut (Germany); Department of Earth and Environmental Sciences, Pusan National University (South Korea); Department of Lithospheric Sciences, Vienna University (Austria) and Mineralogy and Petrology Research Group, University of Granada (Spain). Operating conditions were: accelerating voltage of 15-20 kV, beam current of 10-20 nA, beam size of 3-5 microns.

Trace element mineral chemistry of the xenoliths from the BBHVF was determined by LA-ICP-MS technique using an ELAN 6100 DRC quadrupole mass spectrometer (Institute of Isotope Geochemistry and Mineral Resources, ETH-Zürich, Switzerland) and an ELAN 6100 ICP-MS equipped with CETAC LSX-200 laser ablation system (Geocenter Copenhagen, Denmark), whereas an Agilent 7500ce octopole spectrometer (ORS), whereas GeoLas laser ablation system (Department of Geosciences, Virginia Polytechnic Institute and State University, USA) was used on clinopyroxenes of Jeju Island. The LA-ICP-MS analyses of bulk silicate melt inclusions of the metasomatized peridotites of the BBHVF were carried out using an ELAN 6100 DRC quadrupole mass spectrometer (Institute of Isotope Geochemistry and Mineral Resources, ETH-Zürich, Switzerland). The trace element composition of the coexisting fluid inclusions in these metasomatized peridotites was determined by an Agilent 7500ce octopole spectrometer (ORS) and GeoLas laser ablation system (Department of Geosciences, Virginia Polytechnic Institute and State University, USA).

For microthermometry of fluid inclusions of the peridotites of the BBHVF and Jeju Island, along with the microthermometry of the fluid bubble of silicate melt inclusions in the metasomatized peridotites of the BBHVF a Linkam THMS600 heating-cooling stage mounted on a Nikon Eclipse LV100POL polarizing microscope at Lithosphere Fluid Research Lab (LRG), Eötvös University Budapest (Hungary) was used.

A Jobin Yvon confocal-type Labram Raman instrument with a 532 nm Nd-YAG laser equipped with a CCD detector and 50× objective (Department of Organic Chemistry, Budapest University of Technology and Economics, Hungary) and a similar instrument with a 514 nm laser and 40×

microscope objective (Department of Geosciences, Virginia Polytechnic Institute and State University, USA) were used to analyze fluid composition of the studied peridotites.

The water determination of the minerals in the deformed peridotites of the BBHVF was carried out by Bruker IFS 28 type infrared microscope (Research School of Earth Sciences, The Australian National University, Australia) using unpolarized light and the calibration of Kovács *et al.* (2008) for quantification, whereas infrared analysis on the silicate melt and fluid inclusions along with that of the host minerals in metasomatized peridotites of the BBHVF was performed at ELETTRA Synchrotron Light Laboratory of Trieste (Italy).

## Results

- (1) Based on petrography, deformed spinel lherzolites with flattened tabular equigranular texture and metasomatized amphibole-bearing spinel lherzolites with coexisting silicate melt and fluid inclusions were selected from the BBHVF, whereas three fabric types were distinguished in the Jeju peridotite series forming a continuous transition from the coarse grained-protogranular-porphyroclastic to the fine grained mylonitic textures. Among these rock types, fine grained porphyroclastic and fine grained mylonitic peridotites have not been studied before from the Jeju Island.
- (2) The results of fabric analysis suggest that the deformed peridotites of the BBHVF have extremely strong fabric and olivines possess strong crystallographic preferred orientation with [100]- and [001]-axes homogeneously dispersed in the plane of foliation and [010] showing distinct maximum perpendicular to it. Results of geochemical analyses suggest that these deformed peridotites have enriched trace element composition with spoon-shaped REE distribution. This is a common behavior of equigranular textured peridotites from the BBHVF (Downes *et al.*, 1992) and generally interpreted as a result of cryptic metasomatism of deformed peridotites.
- (3) According to the previous results of deformation analysis on peridotite xenoliths from the central CPR at least two layers can be assumed in the lithospheric mantle having strong deformation patterns and lower equilibrium temperatures at shallower levels and showing the activation of only one slip system with high equilibrium temperatures at the bottom (Falus, 2004). The deformed peridotites studied here do not represent entirely, but might derive from the shallower level and most probably were deformed in a transpressional regime, as suggested based on experimentally deformed rocks. Transpressional regime is suggested to be typical forces acting on the lithosphere in collision or collision-and-escape type geodynamic environments, which was most likely to dominate during the Neogene formation of the Carpathian-Pannonian region, particularly beneath the central Pannonian Basin.
- (4) Based on the trace element and REE patterns of the silicate melt inclusions and their host minerals, as well as the high-temperature crystallization for the clinopyroxene rims, which

enclose petrographically primary inclusions in the metasomatic peridotites of the BBHVF, a mantle-melt interaction can be assumed that occurred at elevated temperature resulted in partial melting of mantle clinopyroxene. This melting was triggered by an evolved reagent melt with trace element content similar to that of the host alkaline basalts, but was richer in SiO<sub>2</sub>.

- (5) Such a melt was likely formed by the interaction of an alkaline mafic melt rising from the asthenosphere with a portion previously metasomatized by probably a slab-derived melt. In the lithospheric mantle after initial clinopyroxene dissolution, the reacting melt likely became clinopyroxene-saturated. Due to cooling, crystallization of the clinopyroxene rims, entrapping drops of melt as silicate melt inclusions, took place. Simultaneously, the microfractures in orthopyroxenes and rarely spinels were filled with the evolved melt, which provided the glassy material (over)saturated in volatiles for the enclosed silicate melt inclusions during fracture healing.
- (6) Coexisting silicate melt and fluid inclusions in the metasomatized peridotites of the BBHVF suggest that both petrographically primary and secondary inclusions were entrapped from an immiscible silicate melt and volatile phase at mantle conditions. The silicate melt inclusions show evolved major element composition with an overall enriched trace element pattern and significant (4-5 wt%) volatile (CO<sub>2</sub>+H<sub>2</sub>O) content. In contrast, the fluid inclusions are C-O-H-S dominated and most probably contain small amounts of silicate components. They possess trace element distribution similar to the silicate melt inclusions, which indicate that a small amount of dissolved silicate melt has a more pronounced effect on the trace element composition of CO<sub>2</sub>-dominated fluid inclusions than the presence of H<sub>2</sub>O. This might be an important conclusion for studies dealing with deep-seated fluid inclusions, especially if they were formed by immiscibility.
- (7) Petrography and major element composition of the peridotite series of Jeju Island show that, apart from the obvious different fabric and grain size distribution; there is no significant difference between the peridotite xenoliths, which derived from the shallow sub-continental lithospheric mantle.
- (8) Conversely, the trace element and REE composition of clinopyroxenes in the Jeju peridotite series indicate that three different chemical groups exist, which form continuous transition from the depleted La<sub>N</sub>/Lu<sub>N</sub> of 0.02 to the spoon-shaped enriched character with La<sub>N</sub>/Lu<sub>N</sub> of 6.37.
- (9) Fabric analysis of the studied peridotites from Jeju Island shed light on the activation of the high temperature (010)[100] slip system, which is responsible for the deformation of the olivines, regardless of texture type. However, in the fine-grained porphyroclastic and fine grained mylonitic peridotites an additional deformation, probably at lower temperature and in a shear-dominated regime significantly weakened the originally strong fabric. This supports the idea that the Jeju peridotites represent a continuous series and suggests that finer the grain size, weaker the fabric.

- (10) Comparing the results of trace element and fabric analysis a strong relationship is revealed between deformation and cryptic metasomatism as the fine grained porphyroclastic and fine grained mylonitic peridotites in every case show enriched trace element character, whereas the coarse grained protogranular-porphyroclastic peridotites are usually depleted. It is not clear whether the shear zones, represented by the fine grained peridotites, encouraged preferential melt infiltration or whether the shear zone is the result of deformation, lubricated by metasomatic agents. Similar conclusion was drawn by Downes (1990), who found relationship between geochemical enrichment and deformation in many mantle peridotites worldwide.
- (11) The composition and evolution of the fluid inclusions in the Jeju peridotite series and in those of the CPR strongly suggest that fluid inclusions trapped at deep lithospheric mantle conditions contain small, thus significant amounts of H<sub>2</sub>O. These observations support our assumption that small amounts of H<sub>2</sub>O exist as a thin film on the walls of many high-density CO<sub>2</sub>-rich fluid inclusions, and its detection is prevented by the inadequacy of currently used analytical techniques. Therefore, the presence of water in deep-seated fluid inclusions is not the special feature of the upper mantle, but rather reflects the general composition of any fluid inclusion from a sub-continental lithospheric setting.

## Acknowledgements

I am grateful to my supervisor, Csaba Szabó (Eötvös University Budapest, Hungary) and my tutors, György Falus and István Kovács (Eötvös Loránd Geophysical Institute, Hungary), as well as Kyoung-hee Yang (Pusan National University, South Korea) for their help. I owe thanks to the members of the Lithosphere Fluid Research Lab (Eötvös University, Budapest) for the fruitful discussions. Special thanks go out to my Family for their patience.

## Published papers and selected abstracts written in the framework of this Ph.D.

### Papers:

- Hidas, K.**, Falus, Gy., Szabó, Cs., Szabó, P.J., Kovács, I. & Földes, T. (2007) Geodynamic implications of flattened tabular equigranular textured peridotites from the Bakony-Balaton Highland Volcanic Field (Western Hungary). *Journal of Geodynamics* 43, 484-503 (Impact Factor: **1.321**, Times Cited: 0)
- Bali, E., Falus, Gy., Szabó, Cs., Peate, D.W., **Hidas, K.**, Török, K. & Ntaflos, T. (2007) Remnants of boninitic melts in the upper mantle beneath the central Pannonian Basin? *Mineralogy and Petrology* 90, 51-72 (Impact Factor: **0.907**, Times Cited: 1)

- Kovács, I., **Hidas, K.**, Hermann, J., Sharygin, V., Szabó, Cs. & Ntaflós, T. (2007) Fluid induced incipient melting in mantle xenoliths from the Minusinsk Region (Siberia, Russia) and some implications for the subcontinental lithospheric mantle. *Geologica Carpathica*, 58, 211-228 (Impact Factor: **0.517**, Times Cited: 0)
- Berkési, M., **Hidas K.**, Szabó, Cs. (2007) Fosszilis geoterma rekonstruálása a Bakony-Balaton-felvidék alatti felsőköpenyben Tihanyról származó peridotit xenolitok CO<sub>2</sub> fluidumzárvány vizsgálatával. *Magyar Geofizika* 48/1, 31-37
- Szabó, Cs., **Hidas, K.**, Bali, E., Zajacz, Z., Kovács, I., Yang, K., Guzmics, T., Török, K. (2009) Melt-wall rock interaction in the mantle shown by silicate melt inclusions in peridotite xenoliths from the central Pannonian Basin (western Hungary). *Island Arc* (in press) (Impact Factor: **0.837**, Times Cited: 0)

#### Abstracts:

- Nam B., Yang K., **Hidas K.**, Um Y., Szabó Cs. (2006) Geochemical features of the subcontinental lithospheric mantle beneath the Jeju Island (South Korea). Annual meeting of the Geological Society of South Korea, November 1-4, 2006, Daejeon (South Korea), Abstract Book
- Hidas, K.**, Yang, K., Berkési, M., Eom, Y., Nam, B. & Szabó, Cs. (2007) Paleo-geotherm reconstruction of the upper mantle: a CO<sub>2</sub> case study from Jeju Island (South Korea). The 5<sup>th</sup> International Symposium on the Volcanoes of the World, May 25-26, 2007, Jeju City, Jeju Island (South Korea), In.: Sung-Hyo Yun (Ed.) Proceedings of the 5<sup>th</sup> international Symposium (JeVI), 27-34
- Nam, B., Yang, K., **Hidas, K.**, Eom, Y., Falus, Gy. & Szabó, Cs. (2007) Foliated peridotites trapped in basaltic rocks from Jeju Island. The 5<sup>th</sup> International Symposium on the Volcanoes of the World, May 25-26, 2007, Jeju City, Jeju Island (South Korea), In.: Sung-Hyo Yun (Ed.) Proceedings of the 5<sup>th</sup> international Symposium (JeVI), 7-13
- Hidas, K.**, Yang, K., Berkési, M., Eom, Y., Nam, B. & Szabó, Cs. (2007) Significance of CO<sub>2</sub> fluid inclusions in the upper mantle: a case study from Jeju Island (South Korea). Proceedings of the Annual Joint Conference, Mineralogical Society of Korea and Petrological Society of Korea. May 31, 2007, Andong, Korea, 35-37
- Nam, B., Yang, K., **Hidas, K.**, Eom, Y., Falus, Gy. & Szabó, Cs. (2007) Shear deformation in the lithospheric mantle beneath Jeju Island, South Korea. (*in Korean*) Proceedings of the Annual Joint Conference, Mineralogical Society of Korea and Petrological Society of Korea. May 31, 2007, Andong, Korea, 90-93
- Szabó, Cs., **Hidas, K.**, Bali, E., Zajacz, Z., Kovács, I., Guzmics, T., Yang, K. & Török, K. (2007) Mafic melt/mantle interaction shown by silicate melt inclusions of peridotite xenoliths from the Bakony-Balaton Highland Volcanic Field (Pannonian Basin, western Hungary).

- Proceedings of the Annual Joint Conference, Mineralogical Society of Korea and Petrological Society of Korea. May 31, 2007, Andong, South Korea, 94-96
- Berkesi, M., **Hidas, K.** & Szabó Cs. (2007) CO<sub>2</sub>-rich fluid inclusions in upper mantle derived xenoliths (Western Hungary, Tihany): a microthermometric and Raman microspectroscopic study. European Current Research on Fluid Inclusions (ECROFI-XIX), 17-20 July, 2007, Bern, Switzerland, Programme and Abstracts, 80
- Hidas, K.**, Szabó, Cs., Guzmics, T., Bali, E., Zajacz, Z. & Kovács, I. (2007) Silicate melt inclusion study in peridotite xenoliths from Pannonian Basin, Hungary. European Current Research on Fluid Inclusions (ECROFI-XIX), 17-20 July, 2007, Bern, Switzerland, Programme and Abstracts, 230
- Hidas, K.**, Szabó, Cs., Guzmics, T., Bali, E., Zajacz, Z. & Kovács, I. (2007) Melt/wallrock interaction shown by silicate melt inclusions in peridotite xenoliths from Pannonian Basin. Goldschmidt 2007 – from atoms to planets, 20-24 August, 2007, Cologne, Germany, Geochimica et Cosmochimica Acta Goldschmidt Abstract Book, A403
- Hidas, K.**, Szabó, Cs., Guzmics, T., Bali, E., Zajacz, Z. & Kovács, I. (2007) Silicate melt inclusions in amphibole-bearing spinel peridotite xenoliths from the Bakony-Balaton Highland Volcanic Field (western Hungary). European Mantle Workshop, Petrological evolution of the European Lithospheric mantle: from Archean to present day, 29-31 August, 2007, Ferrara, Italy, Abstract Volume
- Berkesi, M., **Hidas, K.**, Szabó, Cs. (2007) Pressure preserved by CO<sub>2</sub>-rich fluid inclusions in peridotite xenoliths: reconstructed paleogeotherm within the upper mantle beneath Tihany Peninsula, western Hungary. Annual Meeting of the Hungarian Geological Society, 20-22 September, 2007, Sopron (Hungary), Abstract Book 24-25
- Hidas, K.**, Szabó, Cs., Guzmics, T., Bali, E., Bodnar, R.J., Nédli, Zs. (2008) Significance of H<sub>2</sub>O-bearing, CO<sub>2</sub>-rich fluid inclusions and silicate melt inclusions in upper mantle peridotites from the Bakony-Balaton Highland Volcanic Field. 10<sup>th</sup> Mining, Metallurgy and Geology Conference, April 3-6, 2008, Nagyszeben (Sibiu, Transylvania, Romania), Abstract Book (ISSN 1842-9440)
- Berkesi, M., **Hidas, K.**, Szabó, Cs. (2008) CO<sub>2</sub>-rich fluid inclusion study in upper mantle peridotites (Pannonian Basin, western Hungary). EGU General Assembly, April 16-20, 2008, Vienna (Austria), Abstract Book
- Hidas, K.**, Szabó, Cs., Guzmics, T., Bali E., Bodnar, R.J., Nédli, Zs., Vaccari, L. (2008) C-O-H-S bearing fluids in upper mantle peridotites from the central Pannonian Basin. Pan-American Current Research On Fluid Inclusions (PACROFI-IX), June 2-5, 2008, Washington (USA), Abstract Book
- Hidas, K.**, Szabó, Cs., Guzmics, T., Bali E., Bodnar, R.J., Kovács, I., Nédli, Zs., Perucchi, A., Vaccari, L. (2008) Significance of C-O-H-S bearing fluids and melts in the sub-continental lithospheric mantle of the central Pannonian Basin (W-Hungary). AGU Chapman conference

on shallow mantle composition and dynamics, Fifth international orogenic lherzolite conference, September 22-29, 2008, Mount Shasta (CA, USA), Abstract Book

Szabó Cs., Falus, Gy., Zajacz, Z., Kovács, I., Bali, E., **Hidas, K.** (2008) General geochemical and textural features of the lithospheric mantle beneath the Carpathian-Pannonian region. AGU Chapman conference on shallow mantle composition and dynamics, Fifth international orogenic lherzolite conference, September 22-29, 2008, Mount Shasta (CA, USA), Abstract Book

## References

- Bada, G., Horváth, F., Dövényi, P., Szafián, P., Windhoffer, G. & Cloetingh, S. (2007). Present-day stress field and tectonic inversion in the Pannonian basin. *Global and Planetary Change* 58, 165-180.
- Choi, S. H., Lee, J. I., Park, C. H. & Moutte, J. (2002). Geochemistry of peridotite xenoliths in alkali basalts from Jeju Island, Korea. *Island Arc* 11, 221-235.
- Csontos, L. & Vörös, A. (2004). Mesozoic plate tectonic reconstruction of the Carpathian region. *Palaeogeography Palaeoclimatology Palaeoecology* 210, 1-56.
- Downes, H. (1990). Shear zones in the upper mantle - Relation between geochemical enrichment and deformation in mantle peridotites. *Geology* 18, 374-377.
- Downes, H., Embey-Isztin, A. & Thirlwall, M. F. (1992). Petrology and geochemistry of spinel peridotite xenoliths from the western Pannonian Basin (Hungary) - Evidence for an association between enrichment and texture in the upper mantle. *Contributions to Mineralogy and Petrology* 109, 340-354.
- Falus, G. (2004). Microstructural analysis of upper mantle peridotites: their application in understanding mantle processes during the formation of the Intra-Carpathian Basin System. *Department of Petrology and Geochemistry*. Budapest: Eötvös University, 149.
- Horváth, F., Bada, G., Szafián, P., Tari, G., Ádám, A. & Cloetingh, S. (2006). Formation and deformation of the Pannonian basin: constraints from observational data. In: Gee, D. G. & Stephenson, R. A. (eds.) *European Lithosphere Dynamics*. London: Geological Society, 191-206.
- Kovács, I., Hermann, J., O'Neill, H. S. C., Gerald, J. F., Sambridge, M. & Horváth, G. (2008). Quantitative absorbance spectroscopy with unpolarized light: Part II. Experimental evaluation and development of a protocol for quantitative analysis of mineral IR spectra. *American Mineralogist* 93, 765-778.
- Szabó, C., Falus, G., Zajacz, Z., Kovács, I. & Bali, E. (2004). Composition and evolution of lithosphere beneath the Carpathian-Pannonian Region: a review. *Tectonophysics* 393, 119-137.

