



International Journal of
**Zoological
Research**

ISSN 1811-9778



Academic
Journals Inc.

www.academicjournals.com

Impact of Pesticide Toxicity on Selected Biomarkers in Fishes

Baby Joseph and S. Justin Raj

Interdisciplinary Research Centre, Department of Biotechnology, Malankara Catholic College, Mariagiri, K.K. District, India

Corresponding Author: S. Justin Raj, Interdisciplinary Research Centre, Department of Biotechnology, Malankara Catholic College, Mariagiri, K.K. District, India Tel: 9751358502

ABSTRACT

Modern agricultural practices result in indiscriminate use of various agrochemicals, which usually enter into the aquatic environment. The use of agrochemicals in the field has the potential to change the aquatic medium, affecting the tolerance limit of aquatic fauna and flora, as well as creating danger to the ecosystem. These agrochemicals adversely affect the non-target organisms, especially plankton and fish. The present study reports the acute and sublethal toxicity of pesticides on plasma protein, acetylcholinesterase, hormones, histopathology, changes in gill, ventilatory frequency and stress protein level of freshwater fishes. The alterations of the hormonal levels may be used as a potential biomarker and also can establish the ability of endocrine tissues to respond to their appropriate releasing factors. Heat Shock Proteins (HSPs) are detected in all cells, prokaryotic and eukaryotic. *In vivo* and *in vitro* studies have shown that various stressors transiently increase production of HSPs as protection against harmful insults. Increased levels of HSPs occur after environmental stresses, infection, normal physiological processes and gene transfer.

Key words: Pesticides, hormones, persistency, chlorpyrifos, xenobiotics, stress proteins, HSP70

INTRODUCTION

Pesticides have brought tremendous benefits to mankind by increasing food production and controlling the vectors of man and animal diseases. At the same time use of these pollutants has posed potential health hazards to the life of fishes. Pesticides are major cause of concern for aquatic environment because of their toxicity, persistency and tendency to accumulate in the organisms (Joseph and Raj, 2010). The impact of these pesticides on aquatic organisms is due to the movement of pesticides from various diffuse or point sources. These pesticides are posing a great threat to aquatic fauna especially to fishes, which constitute one of the major sources of protein rich food for mankind (Sharma and Singh, 2007).

BIOCHEMICAL CHANGES IN BLOOD

Blood is highly susceptible to internal and external environment fluctuations because it is the vehicle for the transport of such pollutants (Blaxhall, 1972). Metals are Transported in the blood stream by binding to specific plasma proteins (Joseph *et al.*, 2010). The fish serves as bio-indicator of water quality and the impact of the pesticide can be well understood by analyzing either blood or serum of the fish, because blood is a pathophysiological reflector of whole body (Sharma and Singh, 2004, 2006). The toxic effect of pesticides to the blood of fishes has been

studied by many researchers (Dawson, 1935). The toxicity of chlorpyrifos has been observed on Erythrocyte Sedimentation Rate (ESR) (mm h^{-1}) in fish, *Channa punctatus* (Malla *et al.*, 2009). An increase in ESR (mm h^{-1}) has been reported in *Clarias batrachus* after exposure to savin (Kumar and Benerjee, 1990). Reduction in total serum protein content induces proteinaemia and may be correlated with reduced protein synthesis by liver (Sharma *et al.*, 2009). There was a significant decrease in the total serum protein of cyprinus carpio at 0.1 and 0.001 mL curacron. A significant decrease was observed up to day 21 (Joseph and Raj, 2010). Spontaneous lesions of oocytes can also occur randomly under normal conditions, as reported in zebrafish studies and this is a phenomenon of fish pathology that has been broadly investigated (Rossteuscher *et al.*, 2008). The effects of assimilation of DDT from the water column by fish and this having a mild effect on the parental gonads and direct effects on the F1 generation survival and viability (Mlambo *et al.*, 2009).

ACETYLCHOLINESTERASE

AChE activity is more sensitive for OPs and carbamate pesticides than other contaminants, but the inhibition of this enzyme have been also used to indicate the exposure and effects of other contaminants in fishes. It has been shown that the addition of crude oil to brain homogenate in amounts equivalent to sediment concentration inhibited AChE activity in fishes (Rodriguez-Fuentes and Gold-Bouchot, 2000). Minier *et al.* (2000) reported that muscle AChE of flounder from polluted sites with high level of PAH was inhibited by 40%. Also, a reduction of 40% of brain AChE was observed in *Mullus barbatus* from three pollutes sites of Salento Apulia (Italy), related with presence of great variety of compounds (PAH, heavy metals and pesticides) present in the sediment (Lionetto *et al.*, 2003). The reduction in swimming performance in fish after exposure to Ops could be attributed to the inhibition of AChE (Rao, 2006; Rao *et al.*, 2005). The present results illustrate that after prolonged exposure to PF induced tissue-specific peroxidative damage in brain, gill, viscera and muscle tissues of *O. mossambicus* and the most affected tissue was gill. Earlier studies show that LPO may be induced in various tissues by a variety of environmental pollutants (Ahmad *et al.*, 2000). Significant depression of cholinesterase activities in brain and liver tissues of *O. niloticus* following single and multiple exposure of chlorpyrifos (an organophosphate insecticide) and carbosulfan (a carbamate insecticide) in the laboratory was reported by Chandrasekara and Pathiratne (2005).

HISTOPATHOLOGY

Histopathological changes have been widely used as biomarkers in the evaluation of the health of fish exposed to contaminants, both in the laboratory (Wester and Canton, 1991) and field studies (Hinton *et al.*, 1992; Schwaiger *et al.*, 1997; Teh *et al.*, 1997). One of the great advantages of using histopathological biomarkers in environmental monitoring is that this category of biomarkers allows examining specific target organs, including gills, kidney and liver, that are responsible for vital functions, such as respiration, excretion and the accumulation and biotransformation of xenobiotics in the fish (Gernhofer *et al.*, 2001). Furthermore, the alterations found in these organs are normally easier to identify than functional ones (Fanta *et al.*, 2003) and serve as warning signs of damage to animal health (Hinton and Lauren, 1990).

GILL AS A BIOMARKER

The fish gill is a multifunctional organ involving gaseous exchange acid-base balance, ionic (Na^+ , Cl^- and Ca_2^+) transport and nitrogenous waste excretion (Perry, 1997). The gills consist of

four branchial arches each bearing pairs of primary filaments in which rows of secondary lamellae are situated. The filaments and lamellae are covered by epithelial cells (i.e., pavement cell (PVC), Chloride Cell (CC) and mucous cell) supported by a complex system of blood vessels (Laurent, 1984). Hence, the gill epithelium provides an extensive surface of contact with the environment to facilitate ion transport and gaseous exchange. Gill surfaces are the first target of water-borne metals (Spicer and Weber, 1991). The micro-environment of the gill surface consists of an epithelial membrane which primarily contains phospholipids covered by a mucous layer (Bolis *et al.*, 1984). The constituents of the gill epithelia will be fully ionised, resulting in negatively charged gill surfaces and potential gill-metal interaction sites (Reid and MacDonald, 1991; Joseph and Raj, 2010).

VENTILATORY FREQUENCY

Ventilatory Frequency (VF) represents a good alternative for alert or stressful conditions assessment in fish, because it has several advantages. Investigations have shown that VF is changed quickly in response to disturbances imposed, as demonstrated in fish subjected to social stress (Volpato *et al.*, 1989; Alvarenga and Volpato, 1995), strobe light (Sager *et al.*, 2000), predation risk (Barreto *et al.*, 2003; Hawkins *et al.*, 2004a, b; Queiroz and Magurran, 2005) and confinement (Barreto and Volpato, 2004; Brown *et al.*, 2005). Moreover, VF is an inexpensive and easily measurable parameter, because it can be visually counted, so requiring no sophisticated equipment. It is also a non-invasive behavioural tool, avoiding painful or stressful technique to be used for stress assessment, such as blood sampling.

HORMONES

In fish, hormones are critical towards maintaining proper physiological function and amongst the many hormones found in fish the thyroid hormones (thyroxine (T4) and triiodothyronine (T3)) are known to play an important role in fish growth (Higgs *et al.*, 1982; Miwa and Inui, 1985) and early development (Brown, 1997). When fish are exposed to stressors the levels of thyroid hormones have been demonstrated to be decreased (Pickering, 1993; Deane *et al.*, 2001) and chemical pollutants have been reported to detrimentally affect thyroidal hormone status in a number of fish species (Xu *et al.*, 2002; Brown *et al.*, 2004; Scott and Sloman, 2004; Van der Ven *et al.*, 2006).

STRESS PROTEINS

Exposure of living beings to sub lethal levels of environmental pollution has been shown to trigger several defense mechanisms at the cellular and molecular levels. Very important in this respect is the cellular accumulation of stress proteins which mainly act as molecular chaperones (Ellis and Vander Vies, 1991; Welch, 1993; Bauman *et al.*, 1993; Feder and Hofmann, 1999). Among stress proteins, the HSP70 group (i.e., HSP72, HSP73) has been widely studied since it is regularly over expressed in response to a wide variety of natural, experimental or anthropogenic aggressors viz., heat shock, alcohols, oxidative stress, radiations, heavy metals, arsenic, pesticides and others (Sanders and Martin, 1993; Fulladosa *et al.*, 2002; Gaubin *et al.*, 2002; Delaney and Klesius, 2004). Besides, the stress-related induction of HSP is a general phenomenon observed in all animal species studied so far (Feder and Hofmann, 1999). Metallothioneins (MTs), ubiquitous low molecular weight cysteine-rich proteins, are also considered as stress proteins since they protect the cells against excessive metal uptake (Bauman *et al.*, 1993) by virtue of their high proportion of -SH groups which sequester the metallic ions (Kagi and Schaver, 1988; Klaassen *et al.*, 1999).

As a consequence, MTs should be logically responsive to metal poisoning. Studies carried out in low vertebrates are already numerous and expression of stress proteins in different fish species in response to various stressors has been investigated by many authors (Segner, 1998; Iwama *et al.*, 1998, 1999). For instance, several heat shock proteins have been detected after exposure of various kinds of fish cells to heat shock, arsenate and several metal ions (Kothary and Candido, 1982; Heikkila *et al.*, 1982; Misra *et al.*, 1989; Currie *et al.*, 1999, 2000). Overexpression of MTs has also been studied in different fish species (Carbonell *et al.*, 1998; Dethloff *et al.*, 1999) and its use as a biomarker for monitoring metal pollution in the environment was proposed (Hamilton and Mehrle, 1986). Stress protein mRNA were recently shown to be actively produced in red blood cells of the brook trout submitted to a heat shock (Lund *et al.*, 2003). Besides, fish erythrocytes display high levels of aerobic metabolism that supports many cellular processes, including an active protein synthesis (Currie and Tufts, 1997). Heat shock proteins (hsps), in particular hsp 60 and hsp 70, have been suggested as suitable biomarkers of the exposure to and effects of environmental contaminants (Sanders, 1990). The heat shock proteins are families of proteins which are classified by their molecular weight. They are also known as molecular chaperones (Ellis, 1987) for their constitutive roles in protein synthesis.

HEAT SHOCK PROTEINS

Organisms respond to proteotoxicity with the expression of stress proteins which are able to repair partly denatured proteins. Proteins belonging to the Heat Shock Protein (HSP) families have been demonstrated to play a critical role in the stress response of fish and it has been extensively reported that expression profiles of HSP families can be modulated by abiotic (Iwama *et al.*, 1998; Basu *et al.*, 2002), biotic, heavy metal (Iwama *et al.*, 1998) and organic pollutants. The role of HSP during stress is related to a cytoprotective function as these proteins can act to prevent and repair protein damage (Ananthan *et al.*, 1986). Heat shock protein levels have been shown to be modulated in fish cells and tissues upon exposure to a vast array of stressors (Iwama *et al.*, 1998). Of all the heat shock protein families, HSP70 has been the most comprehensively studied, due to its major importance in cytoprotection (Ryan and Hightower, 1994) and recently it was shown that elevated HSP70 is critical in protection of sea brim cells against chemical induced apoptosis (Deane *et al.*, 2001). The accumulation of these heat shock proteins has been linked to the intensity of stress, these proteins have been regarded as a suitable biomarker in assessing reactions of biota to environmental and physiological stressors (Hightower, 1991; Sanders, 1993).

DISCOVERY OF HSPTS

The first report on HSPs appeared in 1962 after *Drosophila* salivary gland cells were exposed to 37°C for 30 min and then returned to their normal temperature of 25°C for recovery, a puffing of genes was found. All HSP-70s bind ATP (Chappell *et al.*, 1987; Milarski and Morimoto, 1989) have occurred in the chromosome in the recovering cells (Ritossa, 1962) accompanied by an increase in the expression of proteins with molecular masses of 70 and 26 kDa (Tissieres *et al.*, 1974).

CLASSIFICATION OF HSPTS

The best understood HSPs are those with molecular masses of 60, 70, 90 and 110 kDa. These major HSPs are expressed at 37°C in the absence of heat shock. HSP-70 and -90 are observed in all organisms, whereas HSP-110 is present mainly in mammalian cells. A second group of HSPs (sometimes referred as minor HSPs) are induced under conditions of glucose deprivation and include

glucose-regulated proteins (GRP) 34, 47, 56, 75, 78, 94 and 174 kDa (Sciandra and Subject, 1983). A third group of HSPs is the low-molecular-mass HSPs; these have molecular masses of about 20 kDa; these are found at elevated levels in heated *Drosophila* cells (Tissieres *et al.*, 1974) and ischemic cardiomyocytes (Mestril *et al.*, 1994). A small portion of their amino acid sequence is similar to mammalian α -Crystalline (Ingolia and Craig, 1982). HSP-70s are highly conserved and demonstrate a 60-78% base identity among eukaryotic cells and a 40-60% identity between eukaryotic HSP-70 and *Escherichia coli* DnaK, similar to the HSP-70 (Bardwell and Craig, 1984; Craig, 1985; Lindquist, 1986; Caplan *et al.*, 1993).

INDUCTION AND FUNCTIONING OF HEAT SHOCK PROTEINS

Heat shock protein expression also be induced by the presence of denatured proteins (Edington *et al.*, 1989). Increased expression of heat shock proteins has also been called a stress response, because hsp's can be increased or induced after exposure to some environmentally-relevant stressors, including contaminants such as heavy metals (Cd, Cu, Cr, Hg, Ni, Pb and Zn), tributyltin, organophosphate and organochlorine pesticides and other organic contaminants including benzene, 1-chloro-2,4-dinitrobenzene, 2,4-dichloroaniline, 2,4-dinitrobenzene, hexachlorobenzene, pentachlorophenol and trichloroethylene (Sanders, 1993). The common mode of action of these diverse stressors seems to be that they are proteotoxic (Hightower, 1991), resulting in damage to proteins. Basal expression of hsp's has shown to be essential in the early embryonic development and in resting cells. However, over expression of hsp70 in normal drosophila cells slowed down the growth rate, suggesting that over expression of Hsp's in various cells under normal physiological conditions may be of no significance. Molecular chaperones orchestrate when and where proteins fold and unfold in the cell and if there is serious misfolding and aggregation caused by environmental stress or pathology, they can act as sensors to direct these cells to apoptosis. In maintaining the protein homeostasis, Hsp's serve at various levels, such as: (1) protect other proteins against aggregation; (2) prevent protein aggregation; (3) assist the folding of nascent proteins or refolding of damaged proteins; (4) target severely damaged proteins to degradation and (5) in case of excessive damage, sequester damaged proteins to larger aggregates (Soti *et al.*, 2005). The importance of chaperones in promoting and maintaining the native confirmation of cellular proteins is of utmost importance due to the toxic consequences of protein misfolding and aggregation. Cellular accumulation of stress proteins which mainly act as molecular chaperones (Ellis and Vander Vies, 1991; Welch, 1993; Bauman *et al.*, 1993; Feder and Hofmann, 1999). Among stress proteins, the HSP70 group (i.e., HSP72, HSP73) has been widely studied since it is regularly over expressed in response to a wide variety of natural, experimental or anthropogenic aggressors (heat shock, alcohols, oxidative stress, radiations, heavy metals, arsenic, pesticides and others) (Welch, 1993; Sanders and Martin, 1993; Fulladosa *et al.*, 2002; Gaubin *et al.*, 2002; Delaney and Klesius, 2004). Studies carried out in lower vertebrates are already numerous and expression of stress proteins in different fish species in response to various stressors has been investigated by many authors (Segner, 1998; Iwama *et al.*, 1998, 1999). For instance, several heat shock proteins have been detected after exposure of various kinds of fish cells to heat shock, arsenate and several metal ions (Kothary and Candido, 1982; Heikkila *et al.*, 1982; Misra *et al.*, 1989). Protein aggregation associates a number of acute and chronic neurodegenerative conditions. Hsp's reduce the risk of formation of toxic oligomeric assemblies of the respective disease proteins such as tau and amyloid- β in Alzheimers disease, α -synuclein in parkinsons disease huntingtion in Huntingtons disease (Wytttenbach and Arrigo, 2006). Prions are

nothing but infectious particles of protein, which cannot be inactivated by measures that modify nucleic acids. Hsp family members, especially Hsp 140, were shown to inhibit prion protein aggregation independent of ATP. Overexpression of MTs has also been studied in different fish species (Carbonell *et al.*, 1998; Dethloff *et al.*, 1999) and its use as a biomarker for monitoring metal pollution in the environment was proposed (Hamilton and Mehrle, 1986). Stress proteins mRNA were recently shown to be actively produced in red blood cells of the brook trout submitted to a heat shock. Besides, fish erythrocytes display high levels of aerobic metabolism that supports many cellular processes, including an active synthesis of protein. A correlation between levels of heavy metals and metallothioneins in human blood (Szitanyi *et al.*, 1996). The possibility of using MTs gene activation in Peripheral Blood Lymphocytes (PBLs) in humans as a biomarker of susceptibility to cadmium exposure (Lu *et al.*, 2001). Studies carried out on mammalian PBLs of grey seal demonstrated that MTs levels can be induced by Zn (Pillet *et al.*, 2002). However, when using fish as a biological target, only effect of copper in blood and biochemical parameters were analyzed (Dethloff *et al.*, 1999).

STRESS RESPONSE BY HSPS

All HSP-70s bind ATP (Chappell *et al.*, 1987; Milarski and Morimoto, 1989). HSPs are bound to HSFs that reside in the cytosol of mammalian cells under unstressed conditions (Schlesinger, 1990). Under stress conditions such as heat shock and ischemia, HSFs are separated from the HSPs. When HSFs are then phosphorylated by PKC or other serine/threonine kinases, they form a homotrimeric structure (Kroeger *et al.*, 1993). The trimers enter the nucleus, bind to HSEs located on the promoter region of HSP genes and become further phosphorylated by HSF kinases (Price and Calderwood, 1991). Transcription is then initiated, followed by translation. The newly synthesized HSPs bind to HSFs to prevent further synthesis of HSPs (Li *et al.*, 1995).

CONCLUSION

Long term exposure of organisms to pesticides means a continuous health hazard for the population. So, human population is at high risk by consuming these toxicated fishes. This implies that one should take the necessary precaution in the application of pesticides to protect the life of fish and other aquatic fauna. It is known that both environmental and pathological stresses cause an increase in HSPs of host cells. It is believed that levels of HSP-70s might be used as a measure of stress. These organisms can be used to detect the environmental stressors. It is likely that approaches using molecular biology techniques will revolutionize toxicological applications that are cheaper and do not require the use of animals to detect environmental stressors.

REFERENCES

- Ahmad, S., R.K. Scopes, G.N. Rees and B.K.C. Patel, 2000. *Saccharococcus caldoxylosilyticus* sp. nov., an obligately thermophilic, xylose-utilizing, endospore-forming bacterium. *Int. J. Syst. Evol. Microbiol.*, 50: 517-523.
- Alvarenga, C.M.D. and G.L. Volpato, 1995. Agonistic profile and metabolism in alevins of the Nile tilapia. *Physiol. Behav.*, 57: 75-80.
- Ananthan, J., A.L. Goldberg and R. Voellmy, 1986. Abnormal proteins serve as eukaryotic stress signals and trigger the activation of heat shock genes. *Science*, 232: 522-524.
- Bardwell, J.C.A. and E.A. Craig, 1984. Major heat shock gene of drosophila and *Escherichia coli* heat-inducible DNA gene are homologous. *Proc. Natl. Acad. Sci. USA.*, 81: 848-852.

- Barreto, R.E., A.C. Luchiani and A.L. Marcondes, 2003. Ventilatory frequency indicates visual recognition of an allopatric predator in naive Nile tilapia. *Behav. Process.*, 60: 235-239.
- Barreto, R.E. and G.L. Volpato, 2004. Caution for using ventilatory frequency as an indicator of stress in fish. *Behav. Process.*, 66: 43-51.
- Basu, N., A.E. Todgham, P.A. Ackerman, M.R. Bibeau, K. Nakano, P.M. Schulte and G.K. Iwama, 2002. Heat shock protein genes and their functional significance in fish. *Gene*, 295: 173-183.
- Bauman, J.W., J. Liu and C. Klaassen, 1993. Production of metallothioneins and heat shock proteins in response to metals. *Fundamental Applied Toxicol.*, 21: 15-22.
- Blaxhall, P.C., 1972. The haematological assessment of the health of freshwater fish. Review of selected literature. *J. Fish Biol.*, 4: 593-604.
- Bolis, C.L., A. Cambria and M. Fama, 1984. Effects of Acid Stress on Fish Gills. In: *Toxins, Drugs and Pollutants in Marine Mammals*, Bolis, L., J. Zadunaisky and R. Gilles (Eds.). Springer Verlag, Berlin, pp: 122-129.
- Brown, R.M.S., 1997. US assessment of the Georges Bank haddock stock, 1997. The 24th Stock Assessment Workshop/SARC Working Paper Number C1. NEFSC, Woods Hole, MA, pp: 69.
- Brown, S.B., R.E. Evans, L. Vandenbyllardt, K.W. Finnson and V.P. Palace *et al.*, 2004. Altered thyroid status in lake trout (*Salvelinus namaycush*) exposed to co-planer 3, 3', 4, 4', 5-pentachlorobiphenyl. *Aquat. Toxicol.*, 67: 75-85.
- Brown, C., C. Gardner and V.A. Braithwaite, 2005. Differential stress responses in fish from areas of high and low-predation pressure. *J. Comp. Physiol. B*, 175: 305-312.
- Caplan, A.J., D.M. Cyr and M.G. Douglas, 1993. Eukaryotic homologues of *Escherichia coli* dnaJ: A diverse protein family that functions with Hsp70 stress proteins. *Mol. Biol. Cell*, 4: 555-563.
- Carbonell, G., J. Martinez-Pereda and J. Tarazona, 1998. Mobilization of essential metals during and after short-term lethal cadmium exposure in rainbow trout (*Oncorhynchus mykiss*). *Ecotoxicol. Environ. Restor.*, 1: 85-91.
- Chandrasekara, H.U. and A. Pathiratne, 2005. Acetylcholinesterase inhibition and haematological alterations induced in common carp, (*Cyprinus carpio*) following exposure to low concentrations of Trichlorfon. *J. Aquacult. Res.*, 36: 146-150.
- Chappell, T.G., B.B. Konforti, S.L. Schmid and J.E. Rothman, 1987. The ATPase core of a clathrin uncoating protein. *J. Biol. Chem.*, 262: 746-751.
- Craig, E.A., 1985. The heat shock response. *CRC Crit. Rev. Biochem.*, 18: 239-280.
- Currie, S. and B. Tufts, 1997. Synthesis of stress protein 70 Hsp70. in rainbow trout *Oncorhynchus mykiss*. red blood cells. *J. Exp. Biol.*, 200: 607-614.
- Currie, S., B.L. Tufts and C.D. Moyes, 1999. Influence of bioenergetic stress on heat shock protein gene expression in nucleated red blood cells of fish. *Am. J. Physiol.*, 276: R990-R996.
- Currie, S., C.D. Moyes and B.L. Tufts, 2000. The effects of heat shock and acclimation temperature on Hsp70 and Hsp30 mRNA expression in rainbow trout: *In vivo* and *in vitro* comparisons. *J. Fish Biol.*, 56: 398-408.
- Dawson, A.B., 1935. The haematological response in the catfish, *Ameiurus nebulosus* to chronic lead poisoning. *Biol. Bull.*, 68: 335-346.
- Deane, E.E., J. Li and N.Y.S. Woo, 2001. Hormonal status and phagocytic activity in sea bream infected with vibriosis. *Comp. Biochem. Physiol. B*, 129: 687-693.
- Delaney, M.A. and P.H. Klesius, 2004. Hypoxic conditions induce Hsp70 production in blood, brain and head kidney of juvenile Nile tilapia *Oreochromis niloticus* (L.). *Aquacult.*, 236: 633-644.

- Dethloff, G.M., S. Daniel, T.H. Jonathan and C.B. Howard, 1999. Alterations in physiological parameters of rainbow trout (*Onchorynchus mykiss*) with exposure to copper and copper/zinc mixtures. *J. Ecotox. Environ. Safe.*, 42: 253-264.
- Edington, B.V., S.A. Whelan, L.E. Hightower, 1989. Inhibition of heat shock (stress) protein induction by deuterium oxide and glycerol: Additional support for the abnormal protein hypothesis of induction. *J. Cell. Physiol.*, 139: 219-228.
- Ellis, J., 1987. Proteins as molecular chaperones. *Nat.*, 328: 378-379.
- Ellis, R.J. and S.M. Vander Vies, 1991. Molecular chaperones. *Annu. Rev. Biochem.*, 60: 321-347.
- Fanta, E., F.S. Rios, S. Romao, A.C.C. Vianna and S. Freiburger, 2003. Histopathology of the fish *Corydoras paleatus* contaminated with sublethal levels of organophosphorus in water and food. *Ecotoxicol. Environ. Saf.*, 54: 119-130.
- Feder, M.E. and G.E. Hofmann, 1999. Heat shock proteins, molecular chaperones and stress response: Evolutionary and ecological physiology. *Annu. Rev. Physiol.*, 61: 243-282.
- Fulladosa, E., F. Delmas, L. Jun, I. Villaescusa and J.C. Murat, 2002. Cellular stress induced in cultured human cells by exposure to sludge extracts from water treatment plants. *Ecotoxicol. Environ. Safety*, 53: 134-140.
- Gaubin, Y., C. Manceau, D. Skandrani, F. Croute, J.C. Murat and J.P. Soleilhavoup, 2002. Change in level of genes expression after exposure of cultured human cells to endosulfan as investigated by a cDNA microarrays method. *Fresenius Environ. Bull.*, 11: 466-469.
- Gernhofer, M., M. Pawet, M. Schramm, E. Muller and R. Triebkorn, 2001. Ultrastructural biomarkers as tools to characterize the health status of fish in contaminated streams. *J. Aquat. Ecosyst. Stress Recovery*, 8: 241-260.
- Hamilton, S.J. and P.M. Mehrle, 1986. Metallothionein in fish: Review of its importance in assessing stress from metal contaminants. *Trans. Am. Fish. Soc.*, 115: 596-609.
- Hawkins, L.A., A.E. Magurran and J.D. Armstrong, 2004a. Innate predator recognition in newly-hatched Atlantic salmon. *Behav.*, 141: 1249-1262.
- Hawkins, L.A., J.D. Armstrong and A.E. Magurran, 2004b. Predator-induced hyperventilation in wild and hatchery Atlantic salmon fry. *J. Fish Biol.*, 65: 88-100.
- Heikkila, J.J., G.A. Schultz, K. Iatrou and L. Gedamu, 1982. Expression of a set of fish genes following heat or metal ion exposure. *J. Biol. Chem.*, 257: 12000-12005.
- Higgs, D.A., U.H.M. Fagerlund, J.G. Eales and J.R. McBride, 1982. Application of thyroid and steroid hormones as anabolic agents in fish culture. *Comp. Biochem. Physiol. B*, 73: 143-176.
- Hightower, L., 1991. Heat shock, stress proteins, chaperones and proteotoxicity. *Cell*, 66: 191-197.
- Hinton, D.E. and D.J. Lauren, 1990. Liver structural alterations accompanying chronic toxicity in fishes: Potential biomarkers of exposure. In: *Biomarkers of Environmental Contamination*, McCarthy, J.F. and L.R. Shugart (Eds.). Lewis Publishers, Boca Raton FL, USA, ISBN: 0873712846, pp: 51-65.
- Hinton, D.E., P.C. Baumann, G.R. Gardner, W.E. Hawkins and J.D. Hendricks *et al.*, 1992. Histopathologic Biomarkers. In: *Biochemical, Physiological and Histological Markers of Anthropogenic Stress*, Huggett, R.J., R.A. Kimerie, P.M. Mehrie and H.L. Bergman (Eds.). Lewis Publishers, Boac Rato, pp: 155-209.
- Ingolia, T.D. and E.A. Craig, 1982. Four small Drosophila heat shock proteins are related to each other and to mammalian alpha-crystallin. *Proc. Natl. Acad. Sci. USA*, 79: 2360-2364.
- Iwama, G.K., M. Vijayan, R.B. Forsyth and P. Ackerman, 1999. Heat shock proteins and physiological stress in fish. *Am. Zool.*, 39: 901-909.

- Iwama, G.K., P.T. Thomas, R.B. Forsyth and M.M. Vijayan, 1998. Heat shock proteins expression in fish. *Rev. Fish Biol. Fish.*, 8: 35-56.
- Joseph, B. and S.J. Raj, 2010. Effect of curacron toxicity on the total serum protein content of *Cyprinus carpio*. *Toxicol. Environ. Chem.*, 92: 1889-1893.
- Joseph, B., S.J. Raj, T.B. Edwin, P. Sankarganesh, M.V. Jeevitha, S.U. Ajisha and S.R. Sheeja, 2010. Toxic effect of heavy metals on aquatic environment. *Int. J. Biol. Chem. Sci.*, 4: 939-952.
- Kagi, J.H. and A. Schaffer, 1988. Biochemistry of metallothionein. *Biochem.*, 27: 8509-8515.
- Klaassen, C.D., J. Liu and S. Choudhuri, 1999. Metallothionein: An intracellular protein to protect against cadmium toxicity. *Ann. Rev. Pharmacol. Toxicol.*, 39: 267-274.
- Kothary, R.K. and E.P.M. Candido, 1982. Induction of a novel set of polypeptides by heat shock or sodium arsenite in cultured cells of rainbow trout (*Salmo gairdnerii*). *Can. J. Biochem.*, 60: 347-355.
- Kroeger, P.E., K.D. Sarge and R.I. Morimoto, 1993. Mouse heat shock transcription factors 1 and 2 prefer a trimeric binding site but interact differently with the HSP70 heat shock element. *Mol. Cell. Biol.*, 13: 3370-3383.
- Kumar, B. and V. Benerjee, 1990. Effect of sub lethal toxicity of sevin on blood parameters in *Clarias batrachus* (L) Him. *J. Environ. Zool.*, 4: 166-172.
- Laurent, P., 1984. Gill Internal Anatomy. In: *Fish Physiology*, Vol. X A, Hoar, W.S. and D.J. Randall (Eds.). Academic Press, Orlando, pp: 73.
- Li, G.C., S.H. Yang, D. Kim, A. Nussenzweig, H. Ouyang *et al.*, 1995. Suppression of heatinduced hsp70 expression by the 70-kDa subunit of the human Ku autoantigen. *Proc. Natl. Acad. Sci. USA*, 92: 4512-4516.
- Lindquist, S., 1986. The heat-shock response. *Annu. Rev. Biochem.*, 55: 1151-1191.
- Lionetto, M.G., R. Caricato, M.E. Giordano, M.F. Pascariello, L. Marinosci and T. Schettino, 2003. Integrated use of biomarkers (acetylcholinesterase and antioxidant enzymes activities) in *Mytilus galloprovincialis* and *Mullus barbatus* in an Italian coastal marine area. *Mar. Pollut. Bull.*, 46: 324-330.
- Lu, J., T. Jin, G. Nordberg and M. Norberg, 2001. Metallothionein gene expression in peripheral lymphocytes from cadmium-exposed workers. *Cell Stress Chaperones*, 6: 97-104.
- Lund, S., M.E. Lund and B.L. Tufts, 2003. Red blood cell Hsp70 mRNA and protein as bio-indicator of temperature stress in the brook trout (*Salvelinus fontinalis*). *Can. J. Fish. Aquat. Sci.*, 60: 460-470.
- Malla, F.A., G. Sharma and S. Singh, 2009. Chlorpyrifos pesticide toxicity on erythrocyte sedimentation rate in fish, *Channa punctatus* (Bloch.). *Toxeminar-1 Biol. Med.*, 1: 54-55.
- Mestril, R., S.H. Chi, R. Sayen and W.H. Dillman, 1994. Isolation of a novel inducible rat heat-shock protein. HSP70.gene and its expression during ischemiarhypoxia and heat shock. *Biochem. J.*, 298: 561-569.
- Milarski, K.L. and R.I. Morimoto, 1989. Mutational analysis of the human hsp70 protein: Dstinct domains for nucleolar localization and adenosine trisphosphate binding. *J. Cell. Biol.*, 109: 1947-1962.
- Minier, C., F. Levy, D. Rabel, G. Bocquene, D. Godefroy, T. Burgeot and F. Leboulenger, 2000. Flounder health status in the Seine bay. A multim arkers study. *Mar. Environ. Res.*, 50: 373-377.
- Misra, S., M. Zafarullah, J. Price-Haughey and L. Gedamu, 1989. Analysis of stress induced gene expression in fish cell lines exposed to heavy metals and heat shock. *Biochim. Biophys. Acta*, 1007: 325-333.

- Miwa, S. and Y. Inui, 1985. Effects of l-thyroxine and ovine growth hormone on smoltification of amago salmon (*Oncorhynchus rhodurus*). *Gen. Comp. Endocrinol.*, 58: 436-442.
- Mlambo, S.S., J.H.J. van Vuren, I.E.J. Barnhoorn and M.S. Bornman, 2009. Histopathological changes in the reproductive system (ovaries and testes) of *Oreochromis mossambicus* following exposure to DDT. *Environ. Toxicol. Pharmacol.*, 28: 133-139.
- Perry, S.F., 1997. The chloride cell: Structure and functions in the gills of freshwater fishes. *Annu. Rev. Physiol.*, 59: 325-347.
- Pickering, A.D., 1993. Growth and stress in fish production. *Aquacult.*, 111: 51-63.
- Pillet, S., M. Fournier, L.N. Measures, J.M. Bouqueneau and D.G. Cyr, 2002. Presence and regulation of metallothioneins in peripheral blood leukocytes of grey seals. *Toxicol. Applied Pharmacol.*, 185: 207-217.
- Price, B.D. and S.K. Calderwood, 1991. Ca²⁺ is essential for multistep activation of the heat shock factor in permeabilized cells. *Mol. Cell. Biol.*, 11: 3365-3368.
- Queiroz, H. and A.E. Magurran, 2005. Safety in numbers. Shoaling behaviour of the Amazonian red-bellied piranha. *Biol. Lett.*, 1: 155-157.
- Rao, J.V., P. Kavitha, N.M. Jakka, V. Sridhar and P. Usman, 2005. Toxicity of organophosphates on morphology and locomotor behavior in brine shrimp, *Artemia salina*. *Arch. Environ. Contam. Toxicol.*, 539: 227-232.
- Rao, J.V., 2006. Sublethal effects of an organophosphorus insecticide (RPR-II) on biochemical parameters of tilapia, *Oreochromis mossambicus*. *Comp. Biochem. Physiol. C*, 143: 492-498.
- Reid, S.D. and D.G. McDonald, 1991. Metal binding activity of the gills of the rainbow trout (*Oncorhynchus mykiss*). *Can. J. Fish. Aquat. Sci.*, 48: 1061-1068.
- Ritossa, F., 1962. A new puffing pattern induced by temperature shock and DNP in *Drosophila*. *Experientia*, 18: 571-573.
- Rodriguez-Fuentes, G. and G. Gold-Bouchot, 2000. Environmental monitoring using acetylcholinesterase inhibition *in vitro*. A case study in two Mexican lagoons. *Mar. Environ. Res.*, 50: 357-360.
- Rossteuscher, S., H. Schmidt-Posthaus, C. Schaers, M. Teigeler and H. Segner, 2008. Background pathology of the ovary in a laboratory population of zebrafish (*Danio rerio*). *Dis. Aquat. Org.*, 79: 169-172.
- Ryan, J.A. and L.E. Hightower, 1994. Evaluation of heavy metal in fish cells using a combined stress protein and cytotoxicity assay. *Environ. Toxicol. Chem.*, 13: 1231-1240.
- Sager, D.R., C.H. Hocutt and J.R. Stauffer, 2000. Base and stressed ventilation rates for *Leiostomus xanthurus* Lacepede and *Morone Americana* (Gmelin) exposed to strobe lights. *J. Applied Ichthyol.*, 16: 89-97.
- Sanders, B.M., 1990. Stress Proteins: Potential as Multitiered Biomarkers. In: *Biomarkers of Environmental Contamination*, McCarthy, J.F. and L.R. Shugart, (Eds.). Lewis, Boca Raton, pp: 165-191.
- Sanders, B.M., 1993. Stress proteins in aquatic organisms: An environmental perspective. *Crit. Rev. Toxicol.*, 23: 49-75.
- Sanders, B.M. and L.S. Martin, 1993. Stress proteins as biomarkers of contaminant exposure in archived environmental samples. *Sci. Total Environ.*, 139: 459-470.
- Schlesinger, M.J., 1990. Heat shock proteins. *J. Biol. Chem.*, 265: 12111-12114.
- Schwaiger, J., R. Wanke, S. Adam, M. Pawert, W. Honnen and R. Triebkorn, 1997. The use of histopathological indicators to evaluate contaminant-related stress in fish. *J. Aquat. Ecosyst. Stress Recovery*, 6: 75-86.

- Sciandra, J.J. and J.R. Subjeck, 1983. The effects of glucose on protein synthesis and thermosensitivity in Chinese hamster ovary cells. *J. Biol. Chem.*, 258: 12091-12093.
- Scott, G.R. and K.A. Sloman, 2004. The effects of environmental pollutants on complex fish behaviour: Integrating behavioural and physiological indicators of toxicity. *Aquat. Toxicol.*, 68: 369-392.
- Segner, H., 1998. Fish Cell Lines as a Tool in Aquatic Toxicology. In: *Ecotoxicology*, Braunbeck, T., D.E. Hinton and B. Streit (Eds.). Birkhäuser Publishing Ltd., Basel, Switzerland, pp: 1-38.
- Sharma, G. and S. Singh, 2004. Studies on the effect of intoxicant indofil on the blood morphology of *Channa punctatus* (Bloch.). *Bionotes*, 6: 20-20.
- Sharma, G. and S. Singh, 2006. Assay of some blood parameters of the fish, *Channa punctatus* (Bloch.) after intoxication of Indofil. *Bionotes*, 8: 21-21.
- Sharma, G. and S. Singh, 2007. Effect of indofil toxicity on MCHC of *Channa punctatus* (Bloch.). *J. Environ. Res. Dev.*, 1: 261-263.
- Sharma, V., M. Crne, J.O. Park and M. Srinivasarao, 2009. Structural origin of circularly polarized iridescence in jeweled beetles. *Sci.*, 325: 449-451.
- Soti, C., E. Nagy, Z. Giricz, L. Vigh, P. Csermely and P. Ferdinandy, 2005. Heat shock proteins as emerging therapeutic targets. *Br. J. Pharmacol.*, 146: 769-780.
- Spicer, J.I. and R.E. Weber, 1991. Respiratory impairment in crustaceans and molluscs due to exposure to heavy metals. *Comp. Biochem. Physiol. C*, 100: 339-342.
- Szitanyi, Z., C. Nemes and N. Rozlosnik, 1996. Metallothionein and heavy metal concentration in blood. *Microchem. J.*, 54: 246-251.
- Teh, S.J., S.M. Adams and D.E. Hinton, 1997. Histopathological biomarkers in feral freshwater fish populations exposed to different types of contaminant stress. *Aquat. Toxicol.*, 37: 51-70.
- Tissieres, A., H.K. Mitchell and U. Tracy, 1974. Protein synthesis in salivary glands of *Drosophila melanogaster*: Relation to chromosome puffs. *J. Mol. Biol.*, 84: 389-398.
- Van der Ven, L.T.M., E.J. Van den Brandhof, J.H. Vos, D.M. Power and P.W. Wester, 2006. Effects of the antithyroid agent propylthiouracil in a partial life cycle assay with zebrafish. *Environ. Sci. Technol.*, 40: 74-81.
- Volpato, G.L., P.M.A. Frioli and M.P. Carrieri, 1989. Heterogeneous growth in fishes: Some new data in the Nile tilapia, *Oreochromis niloticus* and a general view about the causal mechanisms. *Bol. Fisiol. Anim.*, 13: 7-22.
- Welch, W.J., 1993. How cells respond to stress. *Sci. Am.*, 268: 56-64.
- Wester, P.W. and J.H. Canton, 1991. The usefulness of histopathology in aquatic toxicity studies. *Comp. Biochem. Physiol. C*, 100: 115-117.
- Wytenbach, A. and A.P. Arrigo, 2006. The role of heat shock proteins during neurodegeneration in alzheimer's, parkinson's and huntington's disease. *The Book Heat Shock Proteins in Neural Cells*. Madame Curie Bioscience Database, 1002 West Avenue, Austin, TX 78701. <http://www.ncbi.nlm.nih.gov/bookshelf/br.fcgi?book=eurekah&part=A66297>.
- Xu, Y., J. Zhang, K.W. Schramm and A. Kettrup, 2002. Endocrine effects of sublethal exposure to persistent organic pollutants (POPs) on silver carp (*Hypophthalmichthys molitrix*). *Environ. Pollut.*, 120: 683-690.

The olfactory system is the system related to the sense of smell (olfaction). Many fish activities are dependent on olfaction, such as: mating, discriminating kin, avoiding predators, locating food, contaminant avoidance, imprinting and homing. These activities are referred to as "olfactory-mediated." Impairment of the olfactory system threatens survival and has been used as an ecologically relevant sub-lethal toxicological endpoint for fish within studies. Olfactory information is received by sensory