

## APPENDIX 1B:

### INVENTORY OF HABITAT MODIFICATIONS TO TIDAL INLETS IN THE COASTAL MIGRATION AND WINTERING RANGE OF THE PIPING PLOVER (*Charadrius melodus*)<sup>1</sup>

Tracy Monegan Rice  
Terwilliger Consulting, Inc.  
April 2012

The U.S. Fish and Wildlife Service's (USFWS's) 5-Year Review for the piping plover (*Charadrius melodus*) recommends developing a state-by-state atlas for wintering and migration habitat for the overlapping coastal migration and wintering ranges of the federally listed (endangered) Great Lakes, (threatened) Atlantic Coast and Northern Great Plains piping plover populations (USFWS 2009). The atlas should include data on the abundance, distribution, and condition of currently existing habitat. This assessment addresses this recommendation by providing this data for one habitat type – sandy, tidal inlets within the migration and wintering range of the southeastern continental United States (U.S.). Inlets are a highly valuable habitat for piping plovers and other shorebirds and waterbirds for foraging, loafing, and roosting and have been documented to be preferentially used over other habitat types during the wintering period (Harrington 2008, Lott et al. 2009, Maddock et al. 2009).

While some information is available for the number of inlets stabilized with jetties, revetments, and other hard structures, these data have not been combined with other information that is available for navigational dredging, inlet relocations, shoal mining, and artificial opening and closing of inlets. Altogether this information can provide an assessment of the cumulative impacts of habitat modifications at tidal inlets for piping plovers and other shorebirds and waterbirds. This assessment does **not**, however, include habitat disturbances at tidal inlets such as off-road vehicle (ORV) usage, pet and human disturbance, or disturbance to dunes or vegetation on inlet shoulders.

A description of the different types of stabilization structures typically constructed at or adjacent to inlets – jetties, terminal groins, groins, seawalls, breakwaters and revetments – can be found in Appendix 1A (Rice 2009) as well in the *Manual for Coastal Hazard Mitigation* (Herrington 2003, online at [http://www.state.nj.us/dep/cmp/coastal\\_hazard\\_manual.pdf](http://www.state.nj.us/dep/cmp/coastal_hazard_manual.pdf)) and in *Living by the Rules of the Sea* (Bush et al. 1996).

#### METHODS

This assessment consists chiefly of combining many disparate sources of information regarding tidal inlets within the U.S. continental migration and wintering range into one central Microsoft Excel database. Sources include peer-reviewed literature, books, gray literature (e.g., conference

---


<sup>1</sup> *Suggested citation:*

Rice, T.M. 2012. Inventory of Habitat Modifications to Tidal Inlets in the Coastal Migration and Wintering Range of the Piping Plover (*Charadrius melodus*). Appendix 1B in Draft Comprehensive Conservation Strategy for the Piping Plover (*Charadrius melodus*) Coastal Migration and Wintering Range, U.S. Fish and Wildlife Service. 35 p.

presentations, project applications, or proposals), government reports and files, maps such as Google Earth, U.S. Geological Survey (USGS) topographic maps, nautical charts and state Gazetteers, and on-line databases and government websites (federal, state, county, and municipal).

Google Earth imagery (using the most recent dates available, generally from 2010 and 2011 at inlet locations) and the Federal Inlet Aerial Photo Database (<http://www.oceanscience.net/inletonline/map/map.html>) were utilized to create a database of inlets within the migration and wintering range of the piping plover, namely those within the states of North Carolina, South Carolina, Georgia, Florida, Alabama, Mississippi, Louisiana, and Texas. Zooming in to each inlet allowed identification of existing hard structures and whether the land ownership on the inlet shoulders was developed or undeveloped. Viewing publically-posted digital photographs linked to each location within Google Earth allowed further verification of the existence and type of hard structures or absence thereof.

An inlet is defined as an opening between barrier islands, spits, or peninsulas that allows ocean and bay water to freely exchange and that contains an inlet throat (the main channel) and a series of shoals (Leatherman 1988; Figure 1). Inlets are influenced by sediment supply, the wave climate, the tidal prism (the volume of water passing through the inlet on a tidal cycle), the longshore sediment transport system, sea level rise, and human modifications of the inlet, estuary, river discharging through the inlet, and adjacent shorelines (Leatherman 1988, Davis and Gibeaut 1990, Bush et al. 1996). These various coastal processes and variables are connected with feedback loops, with the inlet's features and behavior seeking to maintain an equilibrium state.



**Figure 1. Schematic diagram of a typical tidal inlet with its morphological features. The ocean or Gulf is to the right in the diagram and the lagoon, bay or estuary is on the left. The net longshore sediment transport is from the top of the diagram to the bottom, the same direction as the dominant waves. Marine waters from the ocean freely exchange with brackish water from the bay, lagoon, sound, or estuary through the inlet on the incoming (flood) and outgoing (ebb) tides. From Schrader et al. (2000).**

Davis and Gibeaut (1990, p. 2) characterize tidal inlets in the following manner:

*Tidal inlets are geologically ephemeral environments which act as dynamic conduits between the sea and coastal bays and which divide the coast into barrier-island segments. Inlets may close and open, migrate or become stable on the order of tens of years in response to changing sediment supply, wave climate and tidal regime, rate of sea level rise, and back-bay filling or dredging. In turn, the associated sediment bodies, ebb- and flood-tidal deltas, may rapidly change character. Because most material making up the inlet sand bodies is taken from the littoral-drift system which feeds adjacent beaches, changes in inlet behavior are reflected by changes in adjacent shorelines and overall barrier-island morphologies .... Tidal inlets are very dynamic and commonly show major changes in inlet size and shape, in some cases even without intervention by man's activities. Changes in wave climate, sediment availability, and nearshore bottom configuration can cause perturbations in coastal processes, and therefore, in the morphology of the inlet or inlets.*

An inlet shoal complex is the group of sand bodies within and near an inlet that is created by an interaction between the tides, waves and sediment supply and consists of both ebb and flood tidal shoals (Figure 1). Individual shoals are separated by tidal channels. Ebb shoals are on the ocean side of an inlet and are more influenced by waves, while flood shoals are on the bay or estuarine side of the inlet and may be emergent during low tide or even maintain some dry (subaerial) lands that could become vegetated over time. A group of ebb tidal shoals is also referred to as an ebb tidal delta, and a group of flood tidal shoals as the flood tidal delta (Leatherman 1988, Bush et al. 1996). Shoals may become relict when an inlet closes, allowing the ebb tidal shoals to weld to the new beach and the flood shoals to stabilize and possibly become vegetated over time. Along deltaic coasts like Louisiana, shoals may become relict if sea level rise outpaces the sediment supply and the inlets essentially drown in place, converting the shoals into subaqueous (submerged) sand bodies and some inlets into open bay mouths. Wide, open bay or sound entrances (e.g., East Cote Blanche Bay in Louisiana, St. George Sound in Florida) were not categorized as inlets in this assessment due to their width and absence of active inlet shoal complexes.

Inlets along deltaic coasts like Louisiana are distinct from tidal inlets typically seen along non-deltaic coasts in the southeastern U.S. The Mississippi and Atchafalaya delta coasts are river-dominated instead of wave- or tide-dominated like other inlets and coasts in the range (Suter 1994). In Texas, the Rio Grande and Brazos River deltas are relatively small and wave-dominated, with most of their distributary streams discharging into estuaries as "bayhead deltas" (Suter 1994, p. 109); as a result, inlets along their Gulf coast segments more resemble tidal inlets along the non-deltaic coasts of the rest of the range (and the flats and shoals associated with the bayhead deltas within the lagoons provide valuable habitat for piping plovers and other birds). Absent human intervention, the deltaic coast of Louisiana would consist of a series of active distributaries, delta plains, extensive wetlands, distributary-mouth bars, and abandoned deltas where marine processes may have reworked the coarser deltaic sediments into barrier islands or spits (transitioning those sections of coast to wave-dominated rather than river-dominated). Abandoned distributary channels may convert into brackish estuaries, as the Bayou Lafourche has done since it ceased to be an active distributary of the Mississippi River roughly 300 years ago (Suter 1994). "The natural geomorphology of a given delta is the result of complex

interactions between sediment supply, relative sea-level changes, and marine reworking. Human interference with any of these factors inevitably alters the form and evolution of the delta. ... [On the Mississippi delta,] through the construction of the levées, the natural processes of the delta were drastically altered. ... Depleted sediment supply from overbanking has accelerated the long-term degradation of the deltaic plain” (Suter 1992, pp. 112-3). As a result, the tidal inlets that are present along the current Louisiana deltaic coast have a very limited sediment supply, preventing them from being self-sustaining without additional sediment input from coastal restoration projects. Where sandy shorelines are present along the Louisiana deltaic coast, tidal inlets were included in this assessment when they exhibited features generally similar to inlets elsewhere in the range (i.e., Figure 1).

Ephemeral breaks or breaches in shorelines or islands were considered inlets in this assessment if they appeared to maintain a tidal exchange of water from the ocean to the bayside; conversely, inlets were considered closed if they did not appear to allow the free flow of water at low tide. This assessment represents a snapshot in time of the inlets open along the southeastern and Gulf coasts of the U.S., using the most recent imagery, publications and personal knowledge available. Inlets are very dynamic, however, and some ephemeral breaches or smaller inlets may have shifted in space or closed and others opened after the publication date of this assessment. Overwash-dominated barrier islands or coasts are especially dynamic, with inlets and breaches repeatedly opening and closing naturally; these areas are included in this survey as a snapshot assessment of the condition of inlet habitats valuable or potentially valuable to the piping plover on its migration and wintering range. The database may be updated in the future with any modifications to the status and new habitat modifications to inlets contained within the geographic area covered in this assessment by contacting the author via email at [tracymrice@yahoo.com](mailto:tracymrice@yahoo.com). Updated copies of the database will be posted on-line at the Program for the Study of Developed Shorelines website (<http://www.wcu.edu/1037.asp>).

Where barrier islands exist offshore of the mainland, entrances or passes that are located on a mainland shoreline are not included as they are geomorphologically distinct from inlets between sandy barrier islands or spits and are estuarine in nature. The mainland coast of Mississippi, for example, provides habitat for the piping plover, but its bay entrances are not included in this assessment since Petit Bois, Horn, Ship and Cat Islands are located offshore and separated by inlets. When the mainland coast does **not** have barrier islands offshore, and the mainland coast is sandy and has direct ocean or Gulf exposure, then mainland passes or inlets are included when they are geomorphically similar to inlets between barrier islands; some of these inlets may be artificially created to provide access to inland water bodies, while others may be river drainages. Examples of these areas include the Matagorda Peninsula in Texas, the Holly Beach area of Louisiana, or the Grand Strand area of South Carolina.

Other mainland areas that do not have sandy coastlines are excluded. The western Florida coast, for example, was limited with one exception to the Northwest Barrier Chain (the Alabama state line east to Ochlockonee Bay in Franklin County) and the West-Central Barrier Chain (Anclote Key south to Cape Romano) and excluded the “plant-dominated, sediment-starved, low-wave energy and tide-dominated coastlines” that are “natural geologic boundaries” of the Big Bend Marsh Coast in between the two barrier island chains and the Ten Thousand Island Mangrove Coast to the south (Hine et al. 2003, p. 2; Davis and Gibeaut 1990, Morton and Peterson 2003a, 2003b, 2004). Bush et al. (2001, p. 171)

characterizes the Big Bend Marsh Coast as an area where “barrier islands are absent and sandy beaches and dunes are rare.” Critical Habitat Unit FL-14 at Hagens Cove in Taylor County, therefore, is excluded from this analysis since it contains no true tidal inlets.

Freshwater or brackish water discharge outlets from lagoons or lakes along the Gulf Coast in the Florida panhandle, Louisiana and Texas were also omitted from this assessment. These outlets generally have no visible tidal deltas, the channels are generally narrow and meandering, and they serve as an outlet for the inland body of water instead of a tidally-flushed inlet between barrier islands or spits. The Florida Keys were also excluded from this inventory, with the area of Atlantic Florida covered stretching from the Georgia state line to Cape Florida on Kckky Biscayne south of Miami Beach. The Florida Keys from Soldier Key south were excluded due to their geologic nature – the islands, or keys, are “a different kind of island chain” that “are quite different from the beach and barrier island systems of East Florida” (Bush et al. 2004, p. 232). The Florida Keys are composed of limestone, are often fringed by mangroves, and natural beaches are quite rare and limited in length (Bush et al. 2004).

Maps in other published sources (e.g., the *Living with the Shore* series of books for individual state coastlines, government reports, journal publications) were then utilized to confirm the number and geographic location of currently open tidal inlets throughout the range, adding non-federally maintained inlet data to the inventory (e.g., inlets dredged by state or local agencies). These map sources were also used to identify the proper political boundaries (i.e., county) in which each inlet is located. News reports and information supplied by relevant public officials and academic sources were consulted to identify the location of new inlets formed within the past few years, typically as a result of storms. History and geology books, literature and government files were referenced to identify inlets that have been relocated or artificially opened or closed since the late 1800s.

In determining the ownership of the inlet shorelines, available maps and on-line directories were searched to identify and verify public properties such as National Wildlife Refuges, National Seashores, state parks and refuges, state wildlife management areas, county and municipal parks and preserves, and lands owned by non-governmental conservation organizations (e.g., Audubon, The Nature Conservancy). Where no records of public ownership were found, the lands were assumed to be privately owned and were recorded as such. Notations were made as to whether the private land was developed or undeveloped; land with low density development such as a small number of structures with no significant infrastructure (e.g., a few fishing cottages) were considered undeveloped due to their dominant land use as being natural.

The primary data source for stabilized inlets was the U.S. Army Corps of Engineers (USACE) Coastal Inlets Research Program (CIRP), which maintains an on-line database of 156 federally-maintained tidal inlets within the U.S. (available at [http://cirp.usace.army.mil/wiki/Inlet\\_Database](http://cirp.usace.army.mil/wiki/Inlet_Database) ). This Federal Inlets Database provides information on stabilization structures including jetties as well as physical characteristics such as tidal prism, inlet dimensions and wave conditions (where data are available); USACE construction history reports are often available for federal structures maintained at inlets included in the database (accessible through

<http://www.oceanscience.net/inletsonline/map/map.html>), providing details on the dates of construction (and thus dates of habitat modification).

These data were combined within a centralized Microsoft Excel database containing the following data fields for each inlet: inlet name, state, north / east land ownership, south / west land ownership, county where the inlet occurs, type of hard structure, location of the structure, structure ownership, date built, dredging (yes or no), dredging maintenance agency, dredged material disposal location(s), sand bypassing (yes or no), shoal mining (yes or no), mining sponsor, date mined, fill location, other miscellaneous but relevant details, and data sources.

A separate Microsoft Excel database was created to catalog the number and location of inlets that have been relocated and naturally and artificially opened and closed since the 1890s. Relocated inlets are those where the inlet is physically moved to a new location – typically hundreds to thousands of feet away – with the old inlet filled in with sediment or other materials and the new inlet excavated through land; inlets generally are relocated as an erosion control method to protect property or infrastructure from loss due to inlet migration. Inlets that were moved to a new location but where the old inlet was not artificially closed and was allowed to remain open were categorized as artificially created inlets and not relocated inlets; if the old inlet subsequently closed naturally, that inlet was categorized as naturally closed. Inlets that have opened or closed due to natural processes include those that were created during storm events or filled in and closed by natural sediment transport processes. Artificially created inlets include those cut through barrier islands or spits where previously no channel existed; these have been created predominantly for navigational purposes but also for water quality or fish passage purposes.

Inlets that have been artificially closed tend to be those opened during a storm event (e.g., Hurricanes Hugo (1989), Katrina (2005) or Irene (2011)) in a location where property owners, governing agencies or politicians consider them undesirable; closure of these new inlets is oftentimes considered a storm recovery endeavor, particularly where it is necessary to restore a road that has been severed by the new inlet. Inlets that have been artificially closed provide a different mosaic of habitats than those that have closed naturally. Inlets that have closed naturally tend to be low, have sparse to no vegetation initially, and wide, especially if the tidal deltas or shoals have welded to the island. Artificially closed inlets, on the other hand, have higher elevations, tend to have a constructed berm and dune system tying in to the adjacent beach and dune systems, and are manually planted with dune grasses and/or other vegetation to stabilize the area; the material used to fill the inlet and construct the berm and dune ridge typically are mined from nearby, disturbing the local sediment supply and transport system. Overwash may occur periodically at a naturally closed inlet but is prevented at an artificially closed inlet by the constructed dune ridge, or in some cases by additional hard structures or sandbags such as those installed at the Rodanthe Breach in North Carolina when it was artificially closed in the fall of 2011. Inlets artificially closed in Louisiana as part of coastal restoration projects are purposefully designed to approximate the natural system and allow overwash in the future, however (B. Firmin, USFWS, personal communication, March 9, 2012). Katrina Cut in Alabama is considered an existing inlet in this assessment (see Table 7) despite its closure with a rock dike during Deepwater Horizon oil spill response

efforts since the dike was permitted as a temporary structure (now undergoing review to remain in place as a permanent structure).

Shoal mining is defined as a project that intentionally mines sediment from a tidal shoal within an inlet complex, typically for beach nourishment purposes on nearby beaches. These projects tend to target the ebb shoals, are located outside of any authorized and/or maintained navigational channels, and tend to require new permits or environmental review. Dredging activities that have occurred within authorized and/or maintained navigational channels with the dredged materials placed on nearby beaches to address erosion are not considered mining projects within this assessment; these types of projects may be considered by the USACE as “beneficial use of dredged material” or as Section 933 projects under the Water Resources Development Act (as amended) but do not create new areas of disturbance to the seafloor as a true mining project does. Both dredging of channels and shoal mining create similar geological and ecological impacts, however, in that they disrupt the sediment transport system within and around inlets, creating sediment sinks within the inlet which can lead to increased erosion rates of adjacent shorelines and sediment bodies.

Data on each inlet were confirmed with multiple sources wherever possible and the sources for each inlet’s data recorded.

The data in both databases were then compiled, sorted and analyzed using common assessment techniques (e.g., the proportion of inlets modified in a particular way within individual states and the range) to identify trends and patterns. Numerous USFWS staff within the range reviewed a draft of the report for verification purposes and corrections / revisions were made accordingly.

## **RESULTS**

Of the 221 tidal inlets that were open in December 2011 within the migration and wintering range of the piping plover, 30 (14%) were artificially created (i.e., cut where there was previously no inlet or dredged open after closing naturally), 8 (4%) have been relocated to new positions entirely, 89 (40%) have been stabilized with one or more hard structures, 97 (44%) have been dredged at least once, and at least 20 (9%) have been mined as a sediment source for beach nourishment. Altogether 119 (54%) of the 221 inlets currently open have been significantly modified in one or more of these manners. Furthermore, at least 64 inlets have been closed artificially and are not included in the 221 total inlets that are presently open (Table 1).

The states with the highest proportion of inlets modified by any means are North Carolina (85%), Atlantic Florida (90%) and the comparably short coast of Alabama, where 100% of the inlets have been modified; in fact, at least 45% of the inlets in all but two states (Georgia and Louisiana) have been modified (Florida has a total of 43 out of 69 inlets modified, or 62%), but that does not take into account the 46 or more inlets in Louisiana that have been artificially closed in recent years (Table 1). Over half (54%) of all the sandy inlets within the migration and wintering range of the piping plover have been modified in one way or another.

Of the 89 inlets that have at least one hard structure, 6 (7%) have one jetty, 45 (49%) have two jetties, 28 (31%) have terminal groins or other groin structures, 24 (27%) have revetments (sandbag or rock) or seawalls, and 4 (4%) have offshore breakwaters (note that the numbers add up to more than 89 since

**Table 1. The number of inlets open as of December 2011 within each state and the migration and wintering range modified by various means.**

State	Existing Inlets							Artificially closed
	Number of Inlets	Total Number of Modified Inlets	Habitat Modification Type					
			structures <sup>†</sup>	dredged	relocated	mined	Artificially opened	
NC	20	17 (85%)	7	16	3	4	2	11
SC	47	21 (45%)	17	11	2	3	0	1
GA	23	6 (26%)	5	3	0	1	0	0
FL – East	21	19 (90%)	19	16	0	3	10	0
FL – West	48	24 (50%)	20	22	0	6	7	1
AL	4	4 (100%)	4	3	0	0	0	2
MS	6	4 (67%)	0	4	0	0	0	0
LA	34	10 (29%)	7	9	1	2	0	46
TX	18	14 (78%)	10	13	2	1	11	3
<b>TOTAL</b>	221	119 (54%)	89 (40%)	97 (44%)	8 (4%)	20 (9%)	30 (14%)	64 (N/A)

<sup>†</sup> Structures include jetties, terminal groins, groin fields, rock or sandbag revetments, seawalls, and offshore breakwaters.

many inlets have more than one type of structure). The highest number of inlets with structures occurs along the Gulf coast of Florida (20) but the highest proportion of inlets stabilized with hard structures occurs along the Atlantic coast of Florida (90%) where only 2 inlets out of 21 have not been stabilized (Table 1).

The state with the highest proportion of natural, unmodified inlets is Georgia (74%). The highest number of contiguous, unmodified, natural inlets is the 15 inlets found between Little Tybee Slough at Little Tybee Island Nature Preserve (GA) and the entrance to Altamaha Sound at the south end of Wolf Island NWR (GA), a distance of approximately 54 miles. The longest stretch of contiguous, unstabilized inlets is located in Louisiana, where 17 inlets between a complex of breaches on the West Belle Pass barrier headland in Lafourche Parish and Beach Prong, located just to the west of the western boundary of the state Rockefeller Wildlife Refuge, have no stabilization structures; one of these inlets, however, is dredged – the Freshwater Bayou Canal. South Carolina also has a lengthy section of coast with no stabilization structures, the 16 inlets from a small unnamed inlet separating the Tom Yawkey Wildlife Center Heritage Preserve from the Santee Coastal Reserve Wildlife Management Area in Georgetown County to Dewees Inlet in Charleston County, although 1 of them has been modified by dredging (Clarks Creek Channel within Bulls Bay). Mississippi is the only state to have no stabilization structures at any of

its 6 inlets, all of which are within Gulf Islands National Seashore where all of the barrier islands are undeveloped (4 of the 6 inlets are dredged, however).

The highest number of inlets that have been modified is along the Atlantic coast of Florida, where 17 of 19 stabilized inlets are contiguous and reach from the St. John's River in Duval County to Norris Cut in Miami-Dade County, a distance of approximately 341 miles; a shorebird would have to travel around 344 miles between unstabilized inlets along this stretch of coast. No other stretch of contiguous inlets longer than 7 inlets (on the Alabama and Florida panhandle coast) has been stabilized within the migration and wintering range of the piping plover, with 3 or 4 contiguous stabilized inlets more common than the extremely high number of contiguous stabilized inlets on the Atlantic coast of Florida.

### **State-specific Results**

#### **North Carolina**

There are 20 tidal inlets currently open in North Carolina, of which 7 (35%) are stabilized with hard structures along at least one shoulder (Table 2). Of the inlets with hard structures, 2 have jetties (one with a single jetty and one with dual jetties), one has a terminal groin, one has a landlocked groin, one has a sandbag groin field, one has a non-functional / submerged breakwater, and 2 have sandbag revetments (one of which also has sheet piling). There are 16 (80%) inlets in North Carolina that have been or continue to be periodically dredged for navigation or erosion control purposes to redirect channels away from buildings or infrastructure. Three inlets (Masonboro Inlet in 1947, Tubbs Inlet in 1970, and Mason Inlet in 2002) have been relocated, with artificial closures of existing inlets and openings of new inlets in a nearby location, while one inlet (Bogue Inlet) had its main channel relocated in 2006 (Masterson et al. 1973, Cleary and Marden 1999, Erickson et al. 2003, Cleary and Fitzgerald 2003, USACE 2004). New inlets have been cut artificially in two locations (Carolina Beach Inlet in 1953, New Drum Inlet in 1971), but neither has been hardened with structures (Pilkey et al. 1998, Mallinson et al. 2008). The shoal complexes of at least 4 inlets have been mined to supply sediment for beach nourishment projects (Shallotte Inlet in 2001, Bogue Inlet in 2005, Barden Inlet in 2006, and Rich Inlet in 1996, 1999 and 2002); two additional inlets have been proposed for mining - Mason Inlet (for Figure Eight Island) and New River Inlet (for Onslow Beach).

At least 11 inlets or breaches have been closed artificially after opening during storm events (Mary's Inlet in the early 1950s, an unnamed breach in Long Beach on Oak Island in 1958, Masonboro Inlet South in 1959, Buxton Inlet in 1963, Moore's Inlet in 1965, Isabel Inlet in 2003, and unnamed breaches on Topsail Island in 1996 and 3 on Hatteras Island near Rodanthe in 2011), while at least 8 inlets were allowed to close as a result of natural coastal processes (New Inlet in 1945, an unnamed inlet on Long Beach in 1956, Mad Inlet in 1997, Old Topsail Inlet in 1998, New / Corncake Inlet in 1999, Old Drum Inlet in 1910, 1971 and 1999, New Drum Inlet in 2008-09, and New-Old Drum Inlet in 2009) (Pilkey et al. 1998, Cleary and Marden 1999, Wamsley and Kraus 2005, Mallinson et al. 2008, Google Earth 2012). Hurricane Isabel in 2003 opened a large new inlet on Hatteras Island near the village of Hatteras, south of Cape Hatteras and within the Cape Hatteras National Seashore, severing North Carolina Highway 12

(NC 12) (Mallinson et al. 2008, Morgan 2009a). The USACE, on behalf of the North Carolina Department of Transportation (NC DOT) and Federal Emergency Management Agency (FEMA) filled in the inlet with

**Table 2. Tidal inlets along the North Carolina coast open as of December 2011 and the types of habitat modification at each. Inlets are presented in geographic order from north to south. Note that an X denotes the modification exists and P denotes it is currently proposed.**

Inlet	Type of Habitat Modification							
	Artificially created	Jetties	Terminal groins / groin field	Seawalls / revetments	Breakwaters	Dredging	Relocation of channel or inlet	Mined for beach fill
Oregon Inlet			X			X		
Pea Island Breach				X				
Hatteras Inlet						X		
Ocracoke Inlet						X		
Ophelia Inlet								
Barden Inlet					X	X		X
Beaufort Inlet		X	X			X		
Bogue Inlet				X		X	X	X
Bear Inlet								
Brown's Inlet								
New River Inlet						X		P
New Topsail Inlet			P			X		
Rich Inlet						X	P	X
Mason Inlet						X	X	P
Masonboro Inlet		X				X	X	
Carolina Beach Inlet	X					X		
Cape Fear River			X			X		
Lockwood's Folly Inlet						X		
Shalotte Inlet						X		X
Tubbs Inlet						X	X	

material dredged from nearby in 40 days, allowing NC DOT to rebuild NC 12 and restore vehicle travel in near record-time (Wamsley and Kraus 2005, Mallinson et al. 2008). Hurricane Irene in August 2011 opened at least 2 inlets and other breaches in and around Rodanthe on Hatteras Island, north of Cape Hatteras and within or adjacent to the Pea Island National Wildlife Refuge (NWR) and Cape Hatteras National Seashore; of these breaches, all but one were filled manually within two months while the most significant new inlet (the Pea Island Breach) was temporarily bridged by the NC DOT while long-term alternatives are evaluated (NC DOT, <http://www.ncdot.gov/travel/nc12recovery/>). On the undeveloped Cape Lookout National Seashore, 2 inlets have opened since 1999 (New-Old Drum and

Ophelia Inlets) and three have naturally closed (Old Drum Inlet, New-Old Drum, and New Drum Inlet – the last of which merged with Ophelia Inlet in 2008-09). Recent studies forecast that the North Carolina Outer Banks will continue to see a series of new inlets open as sea level rises and climate changes (Riggs and Ames 2003, Mallinson et al. 2008).

### South Carolina

There are 47 tidal inlets currently open in South Carolina, of which 17 (36%) are stabilized with hard structures along at least one shoreline (Table 3). Of the inlets with hard structures, 10 have some form of groins (adjacent groins, terminal groins, and/or groin fields), 4 have dual jetties, and 6 have rock revetments and/or seawalls. There are 11 (23%) inlets in South Carolina that have been or continue to be dredged for navigation or erosion control purposes (i.e., to redirect channels away from buildings or

**Table 3. Tidal inlets along the South Carolina coast open as of December 2011 and the types of habitat modification at each. Inlets are presented in geographic order from north to south. Note that an X denotes the modification exists and P denotes it is currently proposed.**

Inlet	Type of Habitat Modification							
	Artificially created	Jetties	Terminal groins / groin field	Seawalls / revetments	Breakwaters	Dredging	Relocation of channel or inlet	Mined for beach fill
Little River Inlet		X				X		
Hog Inlet				X		X		X
Murrell's Inlet		X				X		X
Midway Inlet								
Pawleys Inlet			X					
North Inlet								
Winyah Bay Entrance		X				X		
small unnamed inlet separating Cat or Sand Island from South Island								
North Santee River								
South Santee River								
small unnamed inlet into a lagoon on the north end of Murphy Island adjacent to South Santee River mouth								
Cape Romain Harbor (between Murphy and Cape Islands)								
Unnamed inlet 1 at south end of Cape Island								
Unnamed inlet 2 at south end of Cape Island								

Inlet	Type of Habitat Modification							
	Artificially created	Jetties	Terminal groins / groin field	Seawalls / revetments	Breakwaters	Dredging	Relocation of channel or inlet	Mined for beach fill
Unnamed inlet 3 at south end of Cape Island								
Unnamed inlet separating Cape Island from Lighthouse Island								
Key Inlet								
Unnamed inlet 1 on Raccoon Key								
Unnamed inlet 2 on Raccoon Key								
Bulls Bay						X		
Price Inlet								
Capers Inlet								
Deweese Inlet								
Breach Inlet			X	X				
Charleston Harbor Entrance		X				X		
Lighthouse Inlet			X					
Stono Inlet						X		
small unnamed inlet into tidal lagoon on east end of Kiawah Island						X	X	
Captain Sams Inlet						X	X	
North Edisto River Inlet								
South Creek Inlet								
Frampton Inlet								
Jeremy Inlet				X				
St. Helena Sound Entrance			X					
Johnson Creek			X					
Fripp Inlet			X			X		X
Skull Inlet			X	X				
Price Creek								
Pritchards Inlet								
small unnamed inlet on Little Capers Island								
Trenchards Inlet								
Morse Creek								
Port Royal Sound Entrance			X			X		
Folly Creek			X					
Calibogue Sound Entrance				X				
Mungen Creek			X	X				
Wright River								

infrastructure). One inlet (Captain Sam's Inlet) has been relocated twice (in 1983 and 1996), with artificial closures of the existing inlet and opening of a new inlet in a nearby location (Kana et al. 1987, Lennon et al. 1996). In addition, a project at an unnamed inlet into a tidal lagoon on the east end of Kiawah Island near Stono Inlet relocated the inlet in 2006 and mined material from the adjacent lower beach for use as beach fill on Kiawah Island to the west (USFWS 2006). No new inlets have been artificially created in South Carolina (except for those that were relocated). One inlet or breach on Pawley's Island was closed artificially after opening during Hurricane Hugo in 1989 (Lennon et al. 1996). Eleven new inlets or breaches have opened as a result of storm events in the state since 1989, with Hurricane Irene in August 2011 opening three new breaches on Cape Island at Cape Romain NWR most recently (Lennon et al. 1996, Sarah Dawsey, USFWS Cape Romain NWR pers. Comm.). At least 3 inlets have closed naturally, one in Cherry Grove in the late 1950s (Lennon et al. 1996) and two at Cape Romain NWR around 1992 and 2006 (Sarah Dawsey, USFWS, pers. Comm.). The shoal complexes of at least 3 inlets have been mined to supply sediment for beach nourishment projects (Hog Inlet in 1989/1990, Murrell's Inlet in 1989/1990, and Fripp Inlet in 1975).

## **Georgia**

There are 23 tidal inlets currently open in Georgia, of which 5 (22%) are stabilized with hard structures along at least one shoulder (Table 4). Of the inlets with hard structures, 2 have terminal groins, 2 have adjacent groin fields, 1 has dual jetties, 1 has an offshore breakwater, and 4 have rock revetments and/or seawalls. There are 3 (13%) inlets in Georgia that are dredged for navigation or erosion control purposes. No inlets have been relocated, artificially opened or artificially closed in Georgia. The inlet separating Williamson Island from Little Tybee Island opened naturally sometime between 1957 and 1960 (Clayton et al. 1992), but no other inlets have naturally opened or closed recently in the state. The shoal complex of at least one inlet has been mined to supply sediment for a beach nourishment project (Hampton River Inlet in 1990).

**Table 4. Tidal inlets along the Georgia coast open as of December 2011 and the types of habitat modification at each. Inlets are presented in geographic order from north to south. Note that an X denotes the modification exists and P denotes it is currently proposed.**

Inlet	Type of Habitat Modification							
	Artificially created	Jetties	Terminal groins / groin field	Seawalls / revetments	Breakwaters	Dredging	Relocation of channel or inlet	Mined for beach fill
Savannah River Entrance		X			X	X		
Savannah River South Channel			X	X				
Tybee Inlet			X	X				
Little Tybee Slough								
Little Tybee Creek								
Wassaw Sound Entrance								
Ossabaw Sound Entrance								
Bradley Slough								
Unnamed slough on middle of Ossabaw Island								
Big Slough								
Saint Catherine's Sound Entrance								
Seaside Inlet (at Fish Creek)								
McQueen Inlet								
Sapelo Sound Entrance								
Cabretta Inlet								
Big Hole (between Cabretta and Sapelo Islands)								
Doboy Sound Entrance								
Altamaha Sound Entrance								
Hampton River Inlet						X		X
Gould's Inlet				X				
Saint Simons Sound Entrance				X		X		
Saint Andrews Sound Entrance								
Christmas Creek								

### Florida Atlantic Coast

There are 21 tidal inlets currently open on Florida's Atlantic coast from the Georgia state line south to Key Biscayne, of which 19 (90%) are stabilized with hard structures along at least one shoulder (Table 5). Of the inlets with hard structures, 2 have terminal groins, 16 have jetties (all 16 with 2 jetties), 1 has a rock revetment, 2 have offshore breakwaters and 1 has an adjacent groin field. There are 16 (76%) inlets on Florida's east coast that are dredged for navigation or erosion control purposes (i.e., to redirect

channels away from buildings or infrastructure). No inlets have been relocated, with artificial closures of existing inlets and openings of new inlets in a nearby location. New inlets have been cut artificially in 10 (48%) locations for various purposes (St. Augustine, Sebastian, Fort Pierce, St. Lucie, Lake Worth, Boynton, Boca Raton, Port Everglades, Haulover, and Government Cut Inlets); all of these inlets were cut where no inlets existed at the time except for Boca Raton Inlet, which was repeatedly reopened following natural closures by storms from 1966-1969 and all of them have jetties (Sargent 1988, Bush et al. 2004, Palm Beach County 2003). No inlets have been closed artificially after opening during recent storm events, while four inlets closed as a result of natural coastal processes: old St. Augustine Inlet, Sebastian Inlet in 1941, an inlet near Lake Worth in 1919, and Boca Raton Inlet several times from 1966-1969. Old St. Augustine Inlet between Villano Beach and Conch Island and the one near Black Rocks near Lake Worth were allowed to remain closed, with the other two reopened artificially. A nor'easter in 1973 opened a small breach near Ponce Inlet, which presumably closed shortly thereafter (Bush et al. 2004). An ephemeral inlet periodically opens and closes in the Summer Haven area south of Matanzas Inlet and is currently closed (John Milio, USFWS, pers. Communication 3/8/12). The shoal complexes of at least three inlets have been mined to supply sediment for a beach nourishment project (Boca Raton Inlet in 1985, Jupiter Inlet in 1995, and St. Augustine Inlet in 1996) (Cialone and Stauble 1998, Bush et al. 2004).

**Table 5. Tidal inlets along the east Florida coast open as of December 2011 and the types of habitat modification at each. Inlets are presented in geographic order from north to south. Note that an X denotes the modification exists and P denotes it is currently proposed.**

Inlet	Type of Habitat Modification							
	Artificially created	Jetties	Terminal groins / groin field	Seawalls / revetments	Breakwaters	Dredging	Relocation of channel or inlet	Mined for beach fill
St. Mary's Entrance		X				X		
Nassau Sound Entrance			X					
Fort George Inlet								
St. John's River		X				X		
St. Augustine Inlet	X	X				X		X
Matanzas Inlet				X	X			
Ponce de Leon Inlet		X				X		
Port Canaveral		X				X		
Sebastian Inlet	X	X				X		
Fort Pierce Inlet	X	X				X		
St. Lucie Inlet	X	X			X	X		
Jupiter Inlet		X				X		X
Lake Worth Inlet	X	X				X		
Boynton Inlet (aka South Lake Worth Inlet)	X	X				X		
Boca Raton Inlet	X	X				X		X
Hillsboro Inlet		X				X		
Port Everglades Channel	X	X				X		
Haulover Inlet	X	X	X			X		
Government Cut	X	X				X		
Norris Cut			X					
Bear Cut								

### Florida Gulf Coast

There are 48 tidal inlets currently open along Florida's Gulf coast between the Alabama state line on the panhandle and Cape Romano, of which 20 (42%) are stabilized with hard structures along at least one shoulder (Table 6). Of the inlets with hard structures, 7 have some sort of groin (adjacent groins, terminal groins and/or groin fields), 11 have jetties (3 inlets with 1 jetty and 8 inlets with dual jetties), 1 has an offshore breakwater, and 5 have rock revetments and/or seawalls. There are at least 22 (46%) inlets on Florida's west coast that have been or continue to be dredged periodically for various purposes. No inlets have been relocated along the Gulf coast of Florida, although two new inlets were

opened artificially to replace existing inlets which subsequently closed naturally - new East Pass (Destin Pass) and West Pass (Panama City).

New inlets have been cut artificially (either in new locations or to reopen an inlet that closed naturally) in 8 locations (5 of which now have hard structures): East Pass (Destin Pass) in 1926, West Pass (Saint Andrews Bay - Panama City Harbor) in 1933-1934, Venice Inlet prior to 1937, Bob Sikes Cut in 1954, Clam Pass in 1976 and again in 1981, Midnight Pass in 1983, Blind Pass (Lee County) in 2000 and again in 2009, and St. Andrew Pass on Crooked Island in 2001; a ninth inlet, Big Hickory Inlet, was reopened artificially in 1976 but it closed naturally in 1979 and then reopened naturally since then (Sargent 1988, Davis and Gibeaut 1990, Bush et al. 2001, Antonini et al. 2002). Mexico Beach Canal was also artificially created, is stabilized on both shorelines, and requires dredging to remain open, but it was not included in this assessment since it is a manmade canal with no discernible tidal inlet geomorphology. At least one inlet was closed artificially to block oil spilled in the Deepwater Horizon disaster (Philips Inlet), while at least 17 inlets were allowed to close as a result of natural coastal processes (Sargent 1988, Davis and Gibeaut 1990, Bush et al. 2001, Antonini et al. 2002, Dezember 2010). At least 12 inlets have been opened naturally by storms along the Florida Gulf coast (Sargent 1988, Davis and Gibeaut 1990, Antonini et al. 1999, Bush et al. 2001, Antonini et al. 2002). The shoal complexes of at least 6 inlets have been mined to supply sediment for beach nourishment projects: Pass-a-Grille Channel in the 1980s, Redfish Pass in 1981 and 1988, Johns Pass in 1988, Longboat Pass in 1993, New Pass (Sarasota County) in 1993, and Caxambas Pass in 1990, 1997, 2006 and proposed again for 2012 (Davis and Gibeaut 1990, Cialone and Stauble 1998, Bush et al. 2001, Antonini et al. 2002, Coastal Engineering Consultants 2012). Altogether, 24 of the 48 (50%) west Florida inlets have been modified in some manner.

Table 6. Tidal inlets along the west Florida coast open as of December 2011 and the types of habitat modification at each. Inlets are presented in geographic order from north (west) to south (east) and omit the Big Bend Marsh (except Cedar Keys NWR) and Ten Thousand Island Mangrove sections of coast. Note that an X denotes the modification exists and P denotes it is currently proposed.

Inlet	Type of Habitat Modification							
	Artificially created	Jetties	Terminal groins / groin field	Seawalls / revetments	Breakwaters	Dredging	Relocation of channel or inlet	Mined for beach fill
Pensacola Pass			X			X		
East Pass (aka Destin Pass)	X	X				X		
West Pass (St. Andrew's Bay - Panama City)	X	X				X		
St. Andrew Sound Entrance								
Indian Pass								
West Pass (between St. Vincent and Little St. George Islands)								
Bob Sikes Cut	X	X				X		
East Pass (between eastern St. George Island State Park and Dog Islands)								
Unnamed pass between Anclote Key and Anclote Bar to the north								
Unnamed pass between Three Rooker Bar and Anclote Key								
Unnamed pass between Three Rooker Island and Three Rooker Bar								
St. Joseph Sound (between Honeymoon and Three Rooker Islands)								
Hurricane Pass								
Clearwater Pass		X				X		
Johns Pass		X		X		X		X
Blind Pass (Pinellas County)		X						X
Pass-a-Grille Channel		X				X		X
Bunces Pass								
Unnamed inlet into lagoon on Mullet Key at Fort De Soto Park								
Egmont Channel				X		X		
Southwest Channel								
Passage Key Inlet								
Longboat Pass		X				X		X
New Pass (Sarasota County)			X	X		X		X
Big Sarasota Pass				X				
Venice Inlet	X	X				X		
Stump Pass						X		
Gasparilla Pass								
Boca Grande Pass			X			X		

Inlet	Type of Habitat Modification							
	Artificially created	Jetties	Terminal groins / groin field	Seawalls / revetments	Breakwaters	Dredging	Relocation of channel or inlet	Mined for beach fill
Captiva Pass								
Redfish Pass			X			X		X
Blind Pass (Lee County)	X		X			X		
Matanzas Pass						X		
Big Carlos Pass								
New Pass (Lee County)								
Unnamed breach in Big Hickory Island								
Big Hickory Pass	X <sup>†</sup>		X			X		
Wiggins Pass						X		
Clam Pass	X					X		
Doctors Pass		X				X		
Gordon Pass		X				X		
Little Marco Pass								
Big Marco Pass								
Caxambas Pass			X	X	X	X		X
Unnamed breach between Dickman's and Kice Islands								
Blind Pass (Collier County)								
Unnamed pass between Big Morgan Island and the island to the north								
Morgan Pass								

<sup>†</sup> Big Hickory Inlet closed naturally and was reopened artificially in 1976, but the inlet closed again in 1979; the existing Big Hickory Inlet naturally opened since that time.

## Alabama

There are 4 tidal inlets currently open in Alabama, of which all 4 (100%) are stabilized with hard structures along at least one shoulder (Table 7). Of the inlets with hard structures, one has a groin field, one has dual jetties, one is “temporarily” closed with a rock berm, and 3 have rock or sheet pile revetments and/or seawalls. There are 3 (75%) inlets in Alabama that have been or continue to be dredged periodically. No inlets have been relocated. No new inlets have been cut artificially in Alabama, but West Pass (aka Little Lagoon Pass) was temporarily closed with a sand dike during the Deepwater Horizon oil spill response effort in May 2010 and then was artificially reopened in September 2010 (Dezember 2010). At least two inlets opened on Dauphin Island by hurricanes were allowed to close as a result of natural coastal processes (Bush et al. 2001). Hurricane Ivan in 2004 opened a new inlet at Pine Beach in the Bon Secour NWR (Morgan 2009a), but the inlet appeared closed in 2009

Google Earth imagery. The shoal complexes of no inlets have been mined to supply sediment for beach nourishment projects in Alabama.

Dauphin Island has had several inlets cut across the island by hurricanes, including a 5-mile wide shallow inlet cut by an early 20<sup>th</sup> century hurricane (which was closed by 1942), a September 1948 hurricane, and Hurricane Katrina in 2005 (Bush et al. 2001, USACE 2011). Katrina Cut, opened by Hurricane Katrina on the western end of Dauphin Island, was “temporarily” closed with a rock berm or dike in 2010-2011 with the original purpose to block oil from the Deepwater Horizon oil spill from reaching Mississippi Sound; the state has requested that its temporary permit for the berm from the USACE be converted into a permanent structure (USACE 2011).

**Table 7. Tidal inlets along the Alabama coast open as of December 2011 and the types of habitat modification at each. Inlets are presented in geographic order from west to east. Note that an X denotes the modification exists and P denotes it is currently proposed.**

Inlet	Type of Habitat Modification							
	Artificially created	Jetties	Terminal groins / groin field	Seawalls / revetments	Breakwaters	Dredging	Relocation of channel or inlet	Mined for beach fill
Katrina Cut <sup>†</sup>								
Mobile Pass			X	X		X		
West Pass <sup>‡</sup>				X		X		
Perdido Pass		X		X		X		

<sup>†</sup> Katrina Cut was “temporarily” closed with a rock dike as part of the Deepwater Horizon oil spill response efforts in 2010 but the state is currently seeking permission from the USACE to make the structure permanent (USACE 2011).

<sup>‡</sup> West Pass (aka Little Lagoon Pass) was temporarily closed with a sand dike from in May 2010 as part of the Deepwater Horizon oil spill response efforts but was artificially reopened in September 2010 (Dezember 2010).

## Mississippi

There are 6 tidal inlets currently open in Mississippi, of which none (0%) are stabilized with hard structures along at least one shoulder (Table 8). There are 4 (67%) inlets in Mississippi that are dredged for navigation (Morton 2008). No new inlets have been cut artificially in Mississippi’s barrier islands, been closed artificially after opening during storm events, been relocated, or naturally closed in recent years. At least 7 inlets have been opened by storms in Mississippi, including Camille Cut opened by Hurricane Camille in 1969 on Ship Island, creating West and East Ship Islands (Bowden 1994, Otvos 2006, Otvos and Carter 2008). At least 7 breaches or inlets have closed naturally since 1952 (Otvos and Carter 2008, Stockdon et al. 2010).

The Mississippi Coastal Improvements Program (MsCIP) comprehensive plan for coastal Mississippi proposes the use of dredged material from the Horn Island ship channel to provide beach fill for a portion of West Ship Island within Gulf Islands National Seashore (NPS 2010) and to close Camille Cut between West and East Ship Islands with sediment mined from Sand Island (USACE 2009, Paul Necaie, USFWS, pers. Communication 3/6/12). No other inlet shoal complexes have been mined to supply sediment for beach nourishment projects in Mississippi.

**Table 8. Tidal inlets along the Mississippi coast open as of December 2011 and the types of habitat modification at each. Inlets are presented in geographic order from west to east. Note that an X denotes the modification exists and P denotes it is currently proposed.**

Inlet	Type of Habitat Modification							
	Artificially created	Jetties	Terminal groins / groin field	Seawalls / revetments	Breakwaters	Dredging	Relocation of channel or inlet	Mined for beach fill
Ship Island Pass						X		
Camille Cut <sup>†</sup>								
Dog Keys Pass						X		
Unnamed inlet between Sand Island and Horn Island								
Horn Island Pass						X		
Petit Bois Pass						X		

<sup>†</sup> Camille Cut is proposed to be artificially closed as part of the MsCIP comprehensive plan.

## Louisiana

There are least 34 tidal passes (inlets) with sandy shorelines currently open along the deltaic coast of Louisiana; this total does not include passes without sandy shorelines and counts the Chandeleur Island chain and the West Belle Pass barrier headland as one inlet complex each. The Chandeleur Island chain was fragmented with 44 inlets by Hurricane Katrina in 2005 and restored with roughly 11 inlets closed by the state during the Deepwater Horizon oil spill response effort, resulting in a highly dynamic and uncertain series of islets and inlets. As of September 2011 there were approximately 7 breaches along the West Belle Pass barrier headland shoreline (none of which were present in 2010), but a federally-funded beach restoration project scheduled for 2012 would close any of these breaches that remain open at the time of construction. The vast majority of passes or inlets in Louisiana are connected to extensive wetland complexes and are not inlets separating barrier islands as typically found throughout the rest of the range (and as described in Figure 1); nevertheless, these delta-influenced and sediment-starved inlets often provide valuable shorebird and waterbird habitat.

Of the 34 passes open in 2011, 7 (21%) are stabilized with hard structures along at least one shoreline. Of the inlets with hard structures, 7 have jetties (2 inlets with 1 jetty and 5 inlets with dual jetties), 1 has a groin, and 1 has a rock revetment or seawall (Table 9). There are at least 9 (26%) sandy passes in Louisiana that have been dredged for navigation or other purposes: Calcasieu Pass, Mermentau River, Freshwater Bayou, Belle Pass (Bayou Lafourche), Barataria Pass, Fontanelle Pass, and South Pass of the Mississippi River; Southwest Pass of the Mississippi River is also federally maintained with dredging, but as of 2011 does not have sandy shorelines adjacent to the distributary channel and thus was not included in this analysis. One inlet channel (Bayou Lafourche in 1968) has been relocated, with artificial closure of the existing navigational channel and the opening of a new channel 300 feet to the west (Sargent and Bottin 1989a). No new inlets have been cut artificially (not including oil and gas industry canals).

Breaches cut by Hurricane Andrew (1992) on Raccoon Island were closed artificially (Louisiana Department of Wildlife and Fisheries, <http://www.wlf.louisiana.gov/refuge/terrebonne-barrier-islands-refuge>). Several projects funded under the Coastal Wetlands Planning, Protection and Restoration Act (CWPPRA) have artificially closed inlets while restoring Louisiana's coast, with 3 breaches closed on the Chaland headland in 2006 (CWPPRA Project BA-38), approximately 8 breaches on East Timbalier Island in 1999-2000 (CWPPRA Projects TE-25 and TE-30), 3 breaches on Trinity Island in 1998 (CWPPRA Project TE-24), and the Coupe Nouvelle breach on Whiskey Island in 1998 (CWPPRA Project TE-27) (Louisiana Office of Coastal Protection and Restoration, <http://www.lacoast.gov>). Approximately 29 inlets were closed as part of Deepwater Horizon oil spill response efforts, including 2 on Elmer's Island in Jefferson Parish, approximately 11 in the Chandeleur Island chain, approximately 6 on Scofield Island, approximately 6 on Pelican Island, and approximately 4 on Shell Island as part of the state's sand berms building project (National Commission 2011, Google Earth 2012, Louisiana Office of Coastal Protection and Restoration (<http://coastal.louisiana.gov/index.cfm?md=pagebuilder&tmp=home&pid=131> and <http://www.lacoast.gov>). In addition, any of the 7 breaches on the West Belle Pass barrier headland that are open at the time of construction would be closed as part of a barrier island restoration project funded by the National Oceanic and Atmospheric Administration in October 2011 (NOAA, <http://www.habitat.noaa.gov/hlbarrierislandrestoration.html>). Altogether at least 46 inlets (including those on the Chandeleur Island chain) have been closed artificially in Louisiana in recent years and 7 more are anticipated to be closed in 2012.

Hurricanes Katrina, Rita and others created several dozen new inlets and breaches along the Louisiana coast, most notably within the Chandeleur Island chain of Breton NWR, where the island segmented into 45 islets and 44 inlets/breaches following Hurricane Katrina (Stockdon et al. 2007, Sallenger et al. 2009). An unknown number of inlets have closed as a result of natural coastal processes but the number is likely small as the natural closure of storm breaches during poststorm recovery periods is limited by a restricted supply of sandy sediments in coastal Louisiana and the relatively short period between storms in recent years. At least 2 inlet shoal complexes have been mined to supply sediment for beach nourishment projects (Pass La Mer in 2009 and Chaland Pass in 2009).

**Table 9. Tidal inlets along the Louisiana coast open as of December 2011 and the types of habitat modification at each. Inlets are presented in geographic order from west to east. Note that an X denotes the modification exists and P denotes it is currently proposed; also note that the Chandeleur Island complex is listed here as one entry due to its recent disintegration into dozens of islets, closure of numerous inlets during Deepwater Horizon oil spill response efforts, and uncertain stability.**

Inlet	Type of Habitat Modification							
	Artificially created	Jetties	Terminal groins / groin field	Seawalls / revetments	Breakwaters	Dredging	Relocation of channel or inlet	Mined for beach fill
Unnamed breach in sandbar/spit adjacent to eastern jetty at Sabine Pass								
Calcasieu Pass		X				X		
Mermentau River Navigation Channel		X				X		
Beach Prong (west of Rockefeller Refuge western boundary)								
Joseph Harbor Bayou								
Little Constance Bayou								
Pigeon Bayou								
East Little Constance Bayou								
Rollover Bayou								
Freshwater Bayou Canal						X		
Mosquito Bayou								
Oyster Bayou								
Goreau River								
Bayou de West								
Jack Stout Bayou								
Fish Bayou								
Turtle Bayou								
Whiskey Pass								
Small inlet complex at eastern end of East Timbalier Island								
West Belle Pass barrier headland breaches <sup>†</sup>								
Belle Pass (i.e., Bayou Lafourche)		X				X	X	
Caminada Pass		X						
Barataria Pass		X		X		X		
Pass Abel								
Bayou Quatre Pass								
Pass Ronquille								
Unnamed breach two west of Pass La Mer								
Unnamed breach immediately west of Pass La Mer								
Pass La Mer						X		X
Chaland Pass						X		X
Fontanelle Pass (i.e., Empire Waterway)		X	X			X		

Inlet	Type of Habitat Modification							
	Artificially created	Jetties	Terminal groins / groin field	Seawalls / revetments	Breakwaters	Dredging	Relocation of channel or inlet	Mined for beach fill
Scofield Bayou								
South Pass		X				X		
Chandeleur Island complex								

<sup>†</sup> Any breaches open along the West Belle Pass barrier headland are proposed to be closed in 2012 as part of a federally-funded restoration project (NOAA, <http://www.habitat.noaa.gov/hlbarrierislandrestoration.html>).

## Texas

There are 18 tidal inlets currently open in Texas, of which 10 (56%) are stabilized with hard structures along at least one shoulder (Table 10). Of the inlets with hard structures, 9 have dual jetties, one has groins and one has sheet pile revetments. There are at least 13 (72%) inlets in Texas that have been or continue to be dredged periodically for navigation or other purposes; 8 inlets are federally maintained with navigation channels and 4 have been dredged only one time (Sargent and Bottin 1989b, USACE 1992, Kraus 2007). [Corpus Christi Pass, which is now closed and therefore not included in the above count of 13, was dredged repeatedly in 1928 and 1938 prior to its 1943 closure (the inlet opens and closed intermittently due to storms; USACE 1992)]. The mouth of the Brazos River was relocated 5 miles to the south in 1929 for flood control purposes, but the old river mouth was not closed and currently exists as the Freeport Ship Channel (Sargent and Bottin 1989b, Kraus 2007). The San Bernard River mouth and Bolivar Roads (Galveston Bay) inlets have been relocated (Woody Woodrow, USFWS, pers. Communication 3/6/12). New inlets have been cut artificially in 11 locations for fish passage, flood relief and other purposes in Texas: the Brazos River (Diversion Channel) in 1929, the Colorado River Navigation Channel in 1934, Yarbrough Pass in 1952, Mansfield Pass in 1957 and 1962, Rollover Fish Pass in 1954-55, Matagorda Ship Channel in the 1962, Mustang Island Fish Pass in 1972, McCabe Cut in 1983, Cedar Bayou most recently in 1988 (also in 1939 and 1959), Mitchell's Cut in 1989, and Packery Channel in 2003-06. Four of the artificially created inlets have jetties today. The artificial cuts at both Mustang Island Fish Pass and Yarbrough Pass were unsuccessful and both passages closed naturally, although jetties still exist on the Gulf beach side of Mustang Island Fish Pass (Sargent and Bottin 1989b, USACE 1992, Wamsley and Kraus 2005, Kraus 2007, Williams et al. 2007, Thomas et al. 2011).

Three inlets have been closed artificially – Boca Chica Pass in 1868, Cedar Bayou in 1979 as part of IXTOC oil spill response efforts, and McCabe Cut in 1989 after Mitchell's Cut was opened nearby (USACE 1992). At least 14 inlets have been allowed to close as a result of natural coastal processes: Bryan Beach Cut 2, Wolf Island Cut, Cedar Lakes Pass, Matagorda Peninsula Cut, Brown Cedar Cut, 3-mile Cut, Greens Bayou, Cedar Bayou, Mustang Island Fish Pass, Corpus Christi Pass, Newport Pass, Yarbrough Pass, Mansfield Pass and Boca Chica Pass (Sargent and Bottin 1989b, USACE 1992, Bates 2004, Kraus 2007,

**Table 10. Tidal inlets along the Texas coast open as of December 2011 and the types of habitat modification at each. Inlets are presented in geographic order from west / south to east/ north. Note that an X denotes the modification exists and P denotes it is currently proposed.**

Inlet	Type of Habitat Modification							
	Artificially created	Jetties	Terminal groins / groin field	Seawalls / revetments	Breakwaters	Dredging	Relocation of channel or inlet	Mined for beach fill
Rio Grande River Mouth								
Brazos-Santiago Pass		X	X			X		
Mansfield Pass	X	X				X		
Packery Channel	X	X				X		
Aransas Pass		X				X		
Pass Cavallo						X		
Matagorda Ship Channel	X	X				X		
Colorado River Mouth	X	X				X		
Mitchell's Cut	X					X		
San Bernard River Mouth						X	X	
Brazos River Diversion Channel	X							
Bryan Beach Cut								
Quintana Beach Cut								
Freeport Ship Channel		X				X		
San Luis Pass								P
Bolivar Roads (Galveston Bay)		X				X	X	X
Rollover Pass	X			X		X		X
Sabine Pass		X				X		

Google Earth 2012, Jennifer Wilson, USFWS, pers. Communication 3/7/12). A permit has been issued to reopen Cedar Bayou artificially as a fish pass, but the project has not been constructed yet (Robyn Cobb, USFWS, pers. Communication 3/7/12). At least 5 inlets are hurricane overwash channels that open and close naturally in response to storms, including Brown Cedar Cut, Greens Bayou, Cedar Bayou, Corpus Christi Pass and Newport Pass (USACE 1992, 2003); Cedar Lakes Pass is also a hurricane overwash channel that is influenced by river flows from the San Bernard River but is currently closed (Woody Woodrow, USFWS, pers. Communication 3/7/12). Hurricane Allen reportedly cut 42 breaches across South Padre Island in 1980 (St. John 1991), Hurricane Bret opened a dozen breaches on Padre Island in 1999, and Hurricane Camille opened numerous breaches on Matagorda Island in 1969 (Robyn Cobb, USFWS, pers. Communication 3/7/12). Several overwash breaches appear in Google Earth imagery from 2011 on southern North Padre, South Padre and Brazos Islands from more recent storm events.

Bolivar Roads (Galveston Bay) has been mined to supply sediment for beach nourishment projects and the flood tidal delta of San Luis Pass has been proposed for mining as a source for beach restoration project (Woody Woodrow, USFWS, pers. Communication 3/6/12; Robyn Cobb, USFWS, pers.

Communication 3/7/12). The USACE (1992) reported that 6 of the 8 federally maintained navigation channels contained suitable material for mining as a source for beach fill.

## **DISCUSSION**

Over half (54%) of the sandy, tidal inlet habitat within the U.S. continental migration and wintering range of the piping plover that existed in 2010-2011 has been modified by human actions within the last century or so, including the construction of hard stabilization structures, dredging activities, sediment mining, and the artificial relocation, opening and closing of inlets. The coast of eastern Florida has the most contiguously modified habitat, while significant sections of the South Carolina, Georgia and Louisiana coasts have remained unmodified. Two-thirds or more of the inlets of North Carolina, eastern Florida, Alabama, Mississippi and Texas have been modified (Table 1).

The adverse direct and indirect impacts of hard stabilization structures, dredging, inlet relocations and mining can be significant. The impacts that jetties have on inlet and adjacent shoreline habitat have been described by Cleary and Marden (1999), Bush et al. (1996, 2001, 2004), Wamsley and Kraus (2005), Thomas et al. (2011) and many others. The maintenance of navigation channels by dredging, especially deep ship channels such as those in Alabama and Mississippi, can significantly alter the natural coastal processes on adjacent inlet shorelines and has been described by Otvos (2006), Morton (2008), Otvos and Carter (2008), Beck and Wang (2009), and Stockdon et al. (2010). The relocation of inlets or the creation of new inlets often leads to immediate widening of the new inlet cut and loss of adjacent habitat, amongst other impacts, and has been described by Mason and Sorenson (1971), Masterson et al. (1973), USACE (1992), Cleary and Marden (1999), Cleary and Fitzgerald (2003), Erickson et al. (2003), Kraus et al. (2003), Wamsley and Kraus (2005) and Kraus (2007). Cialone and Stauble (1998) describe the impacts of mining ebb shoals within inlets as a source of beach fill material at 8 locations and provide a recommended monitoring protocol for future mining events; Dabees and Kraus (2008) also describe the impacts of ebb shoal mining.

The cumulative effects of the habitat modifications to sandy tidal inlets within the migration and wintering range of the piping plover are appreciable and significant. The cumulative effects catalogued herein are regional, covering all eight states of the U.S. continental mainland range of the wintering piping plover. Range-wide, over half (54%) of the inlets and their associated habitats have been modified. The cumulative environmental consequences are adverse, major and long-term.

The artificial opening and closing of inlets modifies this type of habitat in the most extreme manner, resulting in the artificial conversion of habitat types and alteration of their abundance and distribution. A high number of inlets (30) have been artificially created within the migration and wintering range of the piping plover, including 10 of the 21 inlets along the eastern Florida coast (Table 1). These artificially created inlets tend to need hard structures to remain open or stable, with 20 of the 30 (67%) of them having hard structures at present. An even higher number of inlets (64) have been artificially closed, the majority in Louisiana; artificial inlet closure results in complete inlet habitat loss. One inlet in Texas was closed as part of the IXTOC oil spill response efforts in 1979, while 32 were closed as part of Deepwater

Horizon oil spill response efforts in 2010-2011. Of the latter, 29 were located in Louisiana, 2 in Alabama and 1 in Florida. To date only one of these inlets, West (Little Lagoon) Pass in Gulf Shores, Alabama, has been reopened, and the rest remain closed with no plans to reopen any of them located during this assessment. The other inlets that have been artificially closed in Louisiana tend to be a result of barrier island restoration projects as much of the state's barrier islands are disintegrating (Otvos 2006, Morton 2008, Otvos and Carter 2008).

The dredging of navigation channels or to relocate inlet channels for erosion control purposes also contributes to the cumulative effects by removing or redistributing the local and regional sediment supply; the maintenance dredging of deep ship channels can convert a natural inlet that normally bypasses sediment from one shoreline to the other into a sediment sink where sediment no longer bypasses the inlet. Of the dredged inlets included in this analysis, dredging efforts began as early as the 1800s and continue to the present, generating long-term and even permanent effects on inlet habitat; at least 11 inlets initiated dredging in the 19<sup>th</sup> century, with the Cape Fear River (NC) dredged as early as 1826 and Mobile Pass (AL) in 1857. Dredging can occur on an annual basis or every 2 to 3 years, resulting in continual perturbations and modifications to inlet and adjacent shoreline habitat. The volumes of sediment removed can be major, with 2.2 million cubic yards (mcy) of sediment removed on average every 1.9 years from the Galveston Bay Entrance (TX) and 3.6 mcy of sediment removed from Sabine Pass (TX) on average every 1.4 years (USACE 1992). The mining of inlet shoals also removes massive amounts of sediment, with 1.98 mcy mined for beach fill from Longboat Pass (FL) in 1998, 1.7 mcy from Shallotte Inlet (NC) in 2001 and 1.6 mcy from Redfish Pass (FL) in 1988 (Cialone and Stauble 1998, USACE 2004). This mining of material from inlet shoals for use as beach fill is not equivalent to the natural sediment bypassing that occurs at unmodified inlets for several reasons, most notably for the massive volumes involved that are "transported" virtually instantaneously instead of gradually and continuously and for the placement of the material outside of the immediate inlet vicinity, where it would naturally bypass. All of the aforementioned dredging and mining impacts are range-wide and occur in every state.

The hard stabilization of inlets is another contributor to the appreciable cumulative effects to inlet habitat along the Gulf of Mexico and southeastern Atlantic coasts of the U.S. The construction of jetties, groins, seawalls and revetments leads to habitat loss and both direct and indirect impacts to adjacent shorelines. Again, these habitat modifications are long-term and permanent, with at least 13 inlets across 6 of the 8 states containing hard structures initially constructed in the 19<sup>th</sup> century. The cumulative effects are on-going and increasing in intensity, with hard structures built as recently as 2011 and others proposed for 2012. With sea level rising and global climate change altering storm dynamics, the pressure to modify the remaining half of sandy, tidal inlets in the range will only increase and would be counterproductive to the adaptation management strategy recommended by the USFWS climate change strategy (USFWS 2010), CCSP (2009), Williams and Gutierrez (2009), Pilkey and Young (2009), and many others.

Indeed, Otvos (2006, p. 1587) found that "[a]ccelerating trends of island destruction have brought delta-fringing Louisiana islands to the verge of extinction." A typical cycle along much of the coast of the migration and wintering range is for storms to open a new inlet or breach in a barrier island, then the

inlet closes naturally as littoral drift slowly fills the breach within a small number of years; in this way islands are segmented, reemerge and are reconstructed as inlets naturally open and close (Davis and Gibeaut 1990, Otvos and Carter 2008, Stockdon et al. 2010). But many sections of coast are disintegrating and in some cases face extinction due to insufficient sediment in the system to support post-storm reconstruction (sometimes due to dredging of nearby channels that act as sediment sinks), more intense and/or frequent storms due to climate change, and a rising sea level, all of which interrupt the cycle. This is presently occurring along the North Carolina Outer Banks (Riggs and Ames 2003, Mallinson et al. 2008, Smith et al. 2008), western Dauphin Island in Alabama (Otvos 2006, Morton 2008), the Mississippi barrier islands of Gulf Islands National Seashore (Morton 2008, Stockdon et al. 2010), and much of the Louisiana coast (Otvos 2006, Morton 2008, Otvos and Carter 2008).

The cumulative effects of the existing habitat modifications to 119 of the 221 inlets as described in this assessment should be addressed in ongoing and future proposals that would affect sandy, tidal inlets within the U.S. continental wintering range of the piping plover. Rising sea level and climate change are likely to increase the number of inlets in the near future. Whether these inlets provide additional favorable habitat to the piping plover and other wildlife resources, however, will depend on the human response to the formation of the new inlets and whether they will be closed, modified in some way, or left to natural processes. The NC DOT and its partners, for example, are currently evaluating long-term solutions to the transportation corridors along the North Carolina Outer Banks and whether to bridge, stabilize or close new inlets such as the ones opened in 2011 by Hurricane Irene. Large-scale plans to restore the Louisiana (Coast 2050 plan) and Mississippi (MsCIP Project) coasts also have been proposed; although these plans would eliminate a significant number of current inlets, they would restore local sediment supplies to maintain beach and inlet habitats and improve their resilience to climate change and rising sea level (Louisiana Coastal Wetlands Conservation and Restoration Task Force and the Wetlands Conservation and Restoration Authority 1998, USACE 2009). Finally, opportunities exist to restore and/or mitigate adverse impacts to existing inlets through the removal of hard structures, elimination of dredging and mining activities, extending dredging cycles to occur less frequently, and the beneficial use of dredged material.

## REFERENCES

- Antonini, G.A., D.A. Fann, and P. Roat. 1999. A Historical Geography of Southwest Florida Waterways. Volume 1 – Anna Maria to Lemon Bay. Florida Sea Grant Program. FLSGP-M-99-004. 80 p. Available at <http://nsgl.gso.uri.edu/flsgp/flsgpm99004/flsgpm99004index.html>.
- Antonini, G.A., D.A. Fann, and P. Roat. 2002. A Historical Geography of Southwest Florida Waterways. Volume 2 – Placida Harbor to Marco Island. Florida Sea Grant Program. FLSGP-M-02-003. Available at <http://nsgl.gso.uri.edu/flsgp/flsgpm02003/flsgpm02003index.html>.
- Barnes, J. 1998. *North Carolina's Hurricane History, Revised and Updated Edition*. Chapel Hill, North Carolina: University of North Carolina Press. 256 p.

- Bates, R.E. 2004. A Brief History of the Laguna Madre Area. Rio Bravo Gallery, Port Isabel, Texas. 2 p. Available at [http://portisabel-texas.com/blog/wp-content/uploads/2010/04/brief\\_history.pdf](http://portisabel-texas.com/blog/wp-content/uploads/2010/04/brief_history.pdf).
- Beck, T.M., and P. Wang. 2009. Influences of channel dredging on flow and sedimentation patterns at microtidal inlets, west-central Florida, USA. *Proceedings of Coastal Dynamics 2009: Impacts of Human Activities on Coastal Processes*, Paper No. 98. Tokyo, Japan. 15 p.
- Bowden, J.E. 1994. *Gulf Islands: The Sands of All Time, Preserving America's Largest National Seashore*. Pensacola, Florida: Pace Printing Company. 143 p.
- Bush, D.M., O.H. Pilkey, Jr., and W.J. Neal. 1996. *Living by the Rules of the Sea*. Durham, North Carolina: Duke University Press. 179 p.
- Bush, D.M., N.L. Longo, W.J. Neal, L.S. Esteves, O.H. Pilkey, D.F. Pilkey, and C.A. Webb. 2001. *Living on the Edge of the Gulf: The West Florida and Alabama Coast*. Durham, North Carolina: Duke University Press. 340 p.
- Bush, D.M., W.J. Neal, N.J. Longo, K.C. Lindeman, D.F. Pilkey, L. Slomp Esteves, J.D. Congleton, and O.H. Pilkey. 2004. *Living with Florida's Atlantic Beaches: Coastal Hazards from Amelia Island to Key West*. Durham, North Carolina: Duke University Press. 338 p.
- Cialone, M.A., and D.K. Stauble. 1998. Historical findings on ebb shoal mining. *Journal of Coastal Research* 14(2):537-563.
- Clayton, T.D., L.A. Taylor, Jr., W.J. Cleary, P.E. Hosier, P.H.F. Graber, W.J. Neal, and O.H. Pilkey, Sr. 1992. *Living with the Georgia Shore*. Durham, North Carolina: Duke University Press. 188 p.
- Cleary, W.J., and D.M. FitzGerald. 2003. Tidal inlet response to natural sedimentation processes and dredging-induced tidal prism changes: Mason Inlet, North Carolina. *Journal of Coastal Research* 19(4):1018-1025.
- Cleary, W.J., and T. Marden. 1999. *Shifting Shorelines: A Pictorial Atlas of North Carolina Inlets*. North Carolina Sea Grant Publication UNC-SG-99-4. 51 p.
- CCSP (Climate Change Science Program). 2009 "Coastal Sensitivity to Sea-Level Rise: A Focus on the Mid-Atlantic Region." A report by the U.S. Climate Change Science Program and the Subcommittee on Global Change Research. [James G. Titus (Coordinating Lead Author), K. Eric Anderson, Donald R. Cahoon, Dean B. Gesch, Stephen K. Gill, Benjamin T. Gutierrez, E. Robert Thieler, and S. Jeffress Williams (Lead Authors)]. U.S. Environmental Protection Agency, Washington D.C., USA. 320 p.
- Coastal Engineering Consultants. 2012. South Marco Renourishment Project Application, Attachment 1: Project Description. January 9, 2012. Submitted to Florida Department of Environmental Protection, Bureau of Beaches and Coastal Systems, Tallahassee, Florida. Available at [http://bcs.dep.state.fl.us/env-prmt/collier/pending/0235209\\_South\\_Marco\\_Island/003\\_JC/Application/Attachment%20I%20-%20Project%20Description/11143Attachment\\_I-20120109.pdf](http://bcs.dep.state.fl.us/env-prmt/collier/pending/0235209_South_Marco_Island/003_JC/Application/Attachment%20I%20-%20Project%20Description/11143Attachment_I-20120109.pdf).

- Coburn, A. S., A. D. Griffith, and R. S. Young, 2010. Inventory of coastal engineering projects in coastal national parks. Natural Resource Technical Report NPS/NRPC/GRD/NRTR—2010/373. National Park Service, Fort Collins, Colorado.
- Dabees, M.A., and N.C. Kraus. 2004. Evaluation of ebb-tidal shoals as a sand source for beach nourishment: General methodology with reservoir model analysis. Proceedings of the 17<sup>th</sup> National Conference on Beach Preservation Technology, Tallahassee, Florida. 21 p. Available at <http://cirp.usace.army.mil/Downloads/PDF/dabees-kraus-FSBPA04.pdf>.
- Dabees, M.A., and N.C. Kraus. 2008. Cumulative effects of channel and ebb shoal dredging on inlet evolution in southwest Florida, USA. Proceedings of the 31<sup>st</sup> International Conference on Coastal Engineering 2008, WS, pp. 2303-2315.
- Davis, R.A., Jr., and J.C. Gibeaut. 1990. Historical Morphodynamics of Inlets in Florida: Models for Coastal Zone Planning. Florida Sea Grant Program, Technical Paper 55. FLSGP-T-90-001 C3. 81 p. Available at <http://nsgl.gso.uri.edu>.
- Davis, R.A., Jr., A.C. Hine, and M.J. Bland. 1987. Midnight Pass, Florida: Inlet instability due to man-related activities in Little Sarasota Bay. Proceedings Coastal Sediments '87, New Orleans, Louisiana. 16 p. Available at <http://nsgl.gso.uri.edu/flsgp/flsgpr87003.pdf>.
- Dean, R.G. 1988. Recommendations for placement of dredged sand on Perdido Key, Gulf Islands National Seashore. Report prepared for the National Park Service, Atlanta, Georgia. University of Florida Publication UFL/COEL-88/016. 51 p. Available at <http://ufdc.ufl.edu/UF00076128/00001/1j>.
- Dezember, R. 2010. "Crews begin reopening Gulf Shores' Little Lagoon Pass." *The Press-Register*, September 10, 2010. Available at [http://blog.al.com/live/2010/09/crews\\_begin\\_reopening\\_gulf\\_sho.html](http://blog.al.com/live/2010/09/crews_begin_reopening_gulf_sho.html).
- Erickson, K.M., N.C. Kraus, and E.E. Carr. 2003. Circulation change and ebb shoal development following relocation of Mason Inlet, North Carolina. *Proceedings Coastal Sediments '03*, World Scientific Publishing Corp. and East Meets West Productions, Corpus Christi, Texas. 13 p.
- Florida Department of Environmental Protection (FL DEP). 1997. Hillsboro Inlet Management Study Implementation Plan, Certificate of Adoption. 7 p. Available at <http://bcs.dep.state.fl.us/bchmngmt/hillsbor.pdf>.
- FL DEP. 2010. Permit Number BA-850 E to the Florida Park Service. Tallahassee, Florida. 12 p. Available at [http://www.dep.state.fl.us/deepwaterhorizon/files/authorizations/071310\\_ba\\_850e.pdf](http://www.dep.state.fl.us/deepwaterhorizon/files/authorizations/071310_ba_850e.pdf).
- Froede, C.R., Jr. 2007. Elevated waves erode the western end of the recently completed sand berm on Dauphin Island, Alabama (U.S.A.). *Journal of Coastal Research* 23(6):1602-1604.
- Google, Inc. 2012. Google Earth (Version 6.2) [Software]. Available from <http://www.google.com/earth/index.html>.

- Harrington, B.R. 2008. Coastal inlets as strategic habitat for shorebirds in the southeastern United States. DOER Technical Notes Collection. ERDC TN-DOER-E25. Vicksburg, Mississippi: U.S. Army Engineer Research and Development Center. Available at <http://el.erd.c.usace.army.mil/dots/doer>.
- Herrington, T.O. 2003. Manual for Coastal Hazard Mitigation. New Jersey Sea Grant College Program, Publication NJSG-03-0511. 108 p. Available at [http://www.state.nj.us/dep/cmp/coastal\\_hazard\\_manual.pdf](http://www.state.nj.us/dep/cmp/coastal_hazard_manual.pdf).
- Hine, A.C., G.R. Brooks, R.A. Davis Jr., D.S. Duncan, S.D. Locker, D.C. Twichell, and G. Gelfenbaum. 2003. The west-central Florida inner shelf and coastal system: A geologic conceptual overview and introduction to the special issue. *Marine Geology* 200(2003):1-17.
- Jaeger, J.M., A. Mehta, R. Faas, and M. Grella. 2009. Anthropogenic impacts on sedimentary sources and processes in a small urbanized subtropical estuary, Florida. *Journal of Coastal Research* 25(1):30-47.
- Kana, T.W., J.E. Mason, and M.L. Williams. 1987. A sediment budget for a relocated tidal inlet. Proceedings Coastal Sediments '87, New Orleans, Louisiana. Abstract available at <http://nsgl.gso.uri.edu/cgi-bin/zgate?present+4932+Default+5+1+F+1.2.840.10003.5.1000.34.1+21>.
- Kraus, N.C. 2007. Coastal Inlets of Texas, USA. *Proceedings Coastal Sediments '07*. Reston, Virginia: ASCE Press. Pp. 1475-1488. Available at <http://www.dtic.mil/cgi-bin/GetTRDoc?Location=U2&doc=GetTRDoc.pdf&AD=ADA481728>.
- Kraus, N.C., G.A. Zarillo, and J.F. Tavolaro. 2003. Hypothetical relocation of Fire Island Inlet, New York. *Proceedings Coastal Sediments '03*. World Scientific Publishing Corp. and East Meets West Productions, Corpus Christi, Texas. Pp. 10-14.
- Lennon, G., W. J. Neal, D.M. Bush, O.H. Pilkey, M. Stutz, and J. Bullock. 1996. *Living with the South Carolina Coast*. Durham, North Carolina: Duke University Press. 241 p.
- Lott, C.A., C.S. Ewell, Jr., and K.L. Volansky. 2009. Habitat associations of shoreline-dependent birds in barrier island ecosystems during fall migration in Lee County, Florida. Dredging Operations and Environmental Research Program Publication ERDC/EL TR-09-14. Engineer Research and Development Center, U.S. Army Corps of Engineers, Washington, D.C. 110 p.
- Louisiana Coastal Wetlands Conservation and Restoration Task Force and the Wetlands Conservation and Restoration Authority. 1998. *Coast 2050: Toward a Sustainable Coastal Louisiana*. Louisiana Department of Natural Resources. Baton Rouge, La. 161 p. Available at <http://www.coast2050.gov/2050reports.htm>.
- Machemehl, J.L., M. Chambers, and N. Bird. 1977. Flow Dynamics and Sediment Movement in Lockwoods Folly Inlet, North Carolina. University of North Carolina Sea Grant Publication UNC-SG-77-11. 153 p. Available at <http://nsgl.gso.uri.edu/ncu/ncut77006.pdf>.

- Maddock, S., M. Bimbi, and W. Golder. 2009. South Carolina shorebird project, draft 2006-2008 piping plover summary report. Audubon North Carolina and U.S. Fish and Wildlife Service, Charleston, South Carolina. 135 pp.
- Mallinson, D.J., S.J. Culver, S.R. Riggs, J.P. Walsh, D. Ames, and C.W. Smith. 2008. Past, Present and Future Inlets of the Outer Banks Barrier Islands, North Carolina. Department of Geological Sciences, East Carolina University, Greenville, North Carolina. 43 p. Available at <http://core.ecu.edu/geology/riggs/Mallinson%20et%20al.%20inlet%20white%20paper%20with%20figures.pdf>.
- Mason, C., and R.M. Sorensen. 1971. Properties and Stability of a Texas Barrier Beach Inlet. Texas A&M University Sea Grant Program Publication No. TAMU-SG-71-217. 177 p. Available at <http://nsgl.gso.uri.edu/tamu/tamut71009.pdf>.
- Masterson, R.P., Jr., J.L. Machemehl, and V.V. Cavaroc. 1973. Sediment Movement in Tubbs Inlet, North Carolina. University of North Carolina Sea Grant Report No. 73-2. 117 p. Available at <http://nsgl.gso.uri.edu/ncu/ncut73013.pdf>.
- McLaughlin, T.E. August 14, 2009. "Homeowners Work to Save Big Hickory Island Beach." *Naples News*. Available at <http://www.naplesnews.com/news/2009/aug/14/videophotos-bonita-homeowners-work-save-beach/>.
- Morgan, K.L.M. 2009a. Coastal Change During Hurricane Isabel 2003. U.S. Geological Survey Fact Sheet 2009-3025. 2 p.
- Morgan, K.L.M. 2009b. Coastal Change During Hurricane Ivan 2004. U.S. Geological Survey Fact Sheet 2009-3026. 2 p.
- Morton, R.A. 2008. Historical changes in the Mississippi-Alabama barrier-island chain and the roles of extreme storms, sea level, and human activities. *Journal of Coastal Research* 24(6):1587-1600.
- Morton, R.A., T. Miller, and L. Moore. 2005. Historical shoreline changes along the US Gulf of Mexico: A summary of recent shoreline comparisons and analyses. *Journal of Coastal Research* 21(4):704-709.
- Morton, R.A., and R.L. Peterson. 2003a. Coastal Classification Atlas: West-Central Florida Coastal Classification Maps - Anclote Key to Venice Inlet. Version 1.1. USGS Open File Report 03-227. Available online with maps at <http://pubs.usgs.gov/of/2003/of03-227/>.
- Morton, R.A., and R.L. Peterson. 2003b. Coastal Classification Atlas: Southwestern Florida Coastal Classification Maps - Venice Inlet to Cape Romano. Version 1.2. USGS Open File Report 03-322. Available online with maps at <http://pubs.usgs.gov/of/2003/of03-322/>.
- Morton, R.A., and R.L. Peterson. 2004. Coastal Classification Atlas: Eastern Panhandle of Florida Coastal Classification Maps - Lighthouse Point to St. Andrew Bay Entrance Channel. USGS Open File Report 2004-1044. Available online with maps at <http://pubs.usgs.gov/of/2004/1044/>.

- National Commission on the BP Deepwater Horizon Oil Spill and Offshore Drilling. 2011. The Story of the Louisiana Berms Project. Staff Working Paper No. 8. 44 p. Available at <http://www.oilspillcommission.gov/sites/default/files/documents/Updated%20Berms%20Working%20Paper.pdf>.
- National Park Service. 2010. Finding of No Significant Impact (FONSI), Proposed West Ship Island North Shore Restoration, Mississippi Sound, Harrison County, Mississippi. U.S. Department of the Interior, National Park Service, Gulf Islands National Seashore. 6 p. Available at <http://www.mscip.usace.army.mil/>.
- North Carolina Department of Environment and Natural Resources (NC DENR). 2011. North Carolina Beach & Inlet Management Plan. Final Report. Raleigh, North Carolina. Various paginations + appendices. Available at <http://dcm2.enr.state.nc.us/BIMP/BIMP%20Final%20Report.html>.
- Otvos, E.G. 2006. Discussion of Froede, C.R., Jr., 2006. The impact that Hurricane Ivan (September 16, 2004) made across Dauphin Island, Alabama. *Journal of Coastal Research*, 22(2), 562-573. *Journal of Coastal Research* 22(6):1585-1588.
- Otvos, E.G., and G.A. Carter. 2008. Hurricane degradation – Barrier island development cycles, northeastern Gulf of Mexico: Landform evolution and island chain history. *Journal of Coastal Research* 24(2):463-478.
- Palm Beach County. 2003. A History of Palm Beach County Inlets. Palm Beach County, Florida. 18 p. Available at <http://www.pbcgov.com>.
- Parchure, T.M. 1982. St. Marys Entrance, Glossary of Inlets Report #11. Florida Sea Grant College Report No. 44. 53 p. Available at <http://nsgl.gso.uri.edu/flsgp/flsgpt82002.pdf>.
- Pilkey, O.H., and R. Young. 2009. *The Rising Sea*. Washington: Island Press. 203 p.
- Pilkey, O.H., W.J. Neal, S.R. Riggs, C.A. Webb, D.M. Bush, D.F. Pilkey, J. Bullock, and B.A. Cowan. 1998. *The North Carolina Shore and Its Barrier Islands: Restless Ribbons of Sand*. Durham, North Carolina: Duke University Press. 318 p.
- Rice, T.M. 2009. Best Management Practices for Shoreline Stabilization to Avoid and Minimize Adverse Environmental Impacts. Prepared for the USFWS, Panama City Ecological Services Field Office. Terwilliger Consulting, Inc., Locustville, Virginia.
- Riggs, S.R. and Ames, D.V., 2003. Drowning the North Carolina coast: Sea-level rise and estuarine dynamics, North Carolina Sea Grant College Program, Raleigh, NC, Pub. No. UNC-SG-03-04, 152 p.
- St. John, B. 1991. *South Padre: The Island and Its People*. Dallas, Texas: Poseidon Press. 281 p.
- Sallenger, A., W. Wright, J. Lillycrop, P. Howd, H. Stockdon, K. Guy, and K. Morgan. 2007. Extreme Changes to Barrier Islands Along the Central Gulf of Mexico Coast During Hurricane Katrina. Chapter 5 in *Science and the Storms: the USGS Response to the Hurricanes of 2005*, G.S. Farris,

- G.J. Smith, M.P. Crane, C.R. Demas, L.L. Robbins, and D.L. Lavois (eds.). U.S. Geological Survey Circular 1306. pp. 113-118.
- Sallenger, A.H., Jr., Wright, C.W., Howd, P., Doran, K., and Guy, K. 2009. Chapter B. Extreme coastal changes on the Chandeleur Islands, Louisiana, during and after Hurricane Katrina. *In* Lavoie, D., ed., Sand resources, regional geology, and coastal processes of the Chandeleur Islands coastal system—an evaluation of the Breton National Wildlife Refuge. U.S. Geological Survey Scientific Investigations Report 2009–5252. pp. 27–36.
- Sargent, F.E. 1988. Case Histories of Corps Breakwater and Jetty Structures. Report 2, South Atlantic Division. Technical Report REMR-CO-3. Waterways Experiment Station, Corps of Engineers, Vicksburg, Mississippi. 105 p.
- Sargent, F.E., and R.R. Bottin, Jr. 1989a. Case Histories of Corps Breakwater and Jetty Structures. Report 8, Lower Mississippi Valley. Technical Report REMR-CO-3. Waterways Experiment Station, Corps of Engineers, Vicksburg, Mississippi. 53 p.
- Sargent, F.E., and R.R. Bottin, Jr. 1989b. Case Histories of Corps Breakwater and Jetty Structures. Report 9, Southwestern Division. Technical Report REMR-CO-3. Waterways Experiment Station, Corps of Engineers, Vicksburg, Mississippi. 53 p.
- Schrader, R.J., M.O. Hayes, T.M. Montello, and E.A. Levine. 2000. Tidal Inlets – A Major Hurdle to Effectively Protecting Sensitive Coastal Resources. *Site Remediation News*, Vol. 12, No. 1, Article 03. Available at [http://www.state.nj.us/dep/srp/news/2000/0007\\_03.htm](http://www.state.nj.us/dep/srp/news/2000/0007_03.htm).
- Smith, C.G., S.J. Culver, S.R. Riggs, D. Ames, D.R. Corbett, and D. Mallinson. 2008. Geospatial analysis of barrier island width of two segments of the Outer Banks, North Carolina, USA: Anthropogenic curtailment of natural self-sustaining processes. *Journal of Coastal Research* 24(1):70-83.
- Stick, D. 1958. *The Outer Banks of North Carolina, 1584-1958*. Chapel Hill, North Carolina: University of North Carolina Press. 352 p.
- Stockdon, H.F., L.A. Fauver, A.H. Sallenger, Jr., and C.W. Wright. 2007. Impacts of Hurricane Rita on the Beaches of Western Louisiana. Chapter 6 in *Science and the Storms: the USGS Response to the Hurricanes of 2005*, G.S. Farris, G.J. Smith, M.P. Crane, C.R. Demas, L.L. Robbins, and D.L. Lavois (eds.). U.S. Geological Survey Circular 1306. pp. 119-123.
- Stockdon, H.F., K.S. Doran, and K.A. Serafin. 2010. Coastal Change on Gulf Islands National Seashore during Hurricane Gustav: West Ship, East Ship, Horn, and Petit Bois Islands. U.S. Geological Survey Open-File Report 2010-1090. 14 p.
- Suter, J.R. 1994. Deltaic Coasts. Chapter 3 in R.W.G. Carter and C.D. Woodroffe (eds.), *Coastal Evolution: Late Quaternary Shoreline Morphodynamics*, Cambridge University Press, pp. 87-120.
- Thomas, R.C., K.B. Brown, and N.C. Kraus. 2011. Inlet stabilization: A case study at Mouth of Colorado River, Texas. *Proceedings of Coastal Sediments 2011*, Miami, Florida. Vol. 1, pp. 533-545.

- United States Army Corps of Engineers (USACE). 1992. Inlets Along the Texas Gulf Coast. Planning Assistance to States Program, Section 22 Report. U.S. Army Engineer District, Galveston, Southwestern Division. 56 p. Available at [http://cirp.usace.army.mil/pubs/archive/Inlets Along TX Gulf Coast.pdf](http://cirp.usace.army.mil/pubs/archive/Inlets_Alone_TX_Gulf_Coast.pdf).
- USACE. 2003. North Padre Island Storm Damage Reduction and Environmental Restoration Project, Nueces County, Texas, Volume 1. Final Environmental Impact Statement. Galveston District, U.S. Corps of Engineers, Galveston, Texas. 297 p. Available at <http://www.swg.usace.army.mil/pe-p/>.
- USACE. 2004. Bogue Inlet Channel Erosion Response Project. Final Environmental Impact Statement. Prepared for The Town of Emerald Isle by Coastal Planning & Engineering, Inc. Wilmington District, U.S. Army Corps of Engineers, Wilmington, North Carolina. Various paginations. Available at <http://www.saw.usace.army.mil/wetlands/Projects/BogueInlet/>.
- USACE. 2009. Comprehensive Plan and Integrated Programmatic Environmental Impact Statement, Mississippi Coastal Improvements Program (MsCIP) Hancock, Harrison, and Jackson Counties, Mississippi. Mobile, Alabama. 417 p. Available at <http://www.ms Cip.usace.army.mil/>.
- USACE. 2011. Public Notice No. SAM-2011-00780-SPG Request for Authorization to Retain and Maintain and Existing Sand-filled Rock Rubble Berm, West end of Dauphin Island, Gulf of Mexico/Mississippi Sound, Mobile County, Alabama. Mobile District, Corps of Engineers, Mobile, Alabama. 26 p.
- U.S. Fish and Wildlife Service (USFWS). 2006. Biological Opinion for the Kiawah Island Beach Nourishment Project, Charleston County, South Carolina. Charleston Ecological Services Field Office, Charleston, South Carolina. 59 p.
- USFWS. 2009. Piping plover (*Charadrius melodus*) 5-year review: summary and evaluation. Northeast Region, Hadley, Massachusetts. 206 p.
- USFWS. 2010. Rising to the Urgent Challenge: Strategic Plan for Responding to Accelerating Climate Change. Washington, D.C. 32 p.
- Wamsley, T.V., and N.C. Kraus. 2005. Coastal Barrier Island Breaching, Part 2: Mechanical Breaching and Breach Closure. U.S. Army Corps of Engineers Technical Note ERDC/CHL CHETN-IV-65. 21 p.
- Williams, S.J., and B. Gutierrez. 2009. Sea-level rise and coastal change: Causes and implications for the future of coasts and low-lying regions. *Shore and Beach* 77(4):13-21.
- Williams, D.D., N.C. Kraus, and C.M. Anderson. 2007. Morphologic response to a new inlet, Packery Channel, Corpus Christi, Texas. *Proceedings Coastal Sediments '07 Conference*, ASCE Press, Reston, VA. Pp. 1529-1542.

Terwilliger Consulting Inc., Locustville, VA. 190 likes. TCI offers a full range of professional natural resource/ wildlife/environmental conservation. See more of Terwilliger Consulting Inc. on Facebook. Log In. or. Create New Account. See more of Terwilliger Consulting Inc. on Facebook. Log In. Forgot account? Contact. More. © 2016 by Terwilliger Consulting Inc. PROJECTS. More Info. Terwilliger can refer to: Brian J. Terwilliger, movie director. George J. Terwilliger III, former US Deputy Attorney General. George Terwilliger, American silent film director and screenwriter. Joseph Terwilliger, American geneticist. Robert Underdunk "Sideshow Bob" Terwilliger, a fictional character from The Simpsons. Wayne Terwilliger, former second baseman in Major League Baseball. United States: Terwilliger Boulevard, a major road and parkway in Portland, Oregon.