

Original Research Article

doi: <http://dx.doi.org/10.20546/ijcmas.2016.502.031>

Microbial Vaginitis in Reproductive Women of Silk city of South India

Karthika Jayakumar^{1*}, Wills shiela², S. Jayalakshmi¹,
Sathya Pandurangan¹ and M. Mohanambal²

¹Department of Microbiology, Shri Sathya Sai Medical College and Research Institute
Kancheepuram (District), Tamilnadu, India

²Department of O and G, Shri Sathya Sai Medical College and Research Institute,
Kancheepuram (District), Tamilnadu, India

*Corresponding author

ABSTRACT

Keywords

Trichomonas vaginalis,
Culture,
Microscopy,
Vaginitis,
Silk city of
South India

Article Info

Accepted:
15 January 2016
Available Online:
10, February 2016

Vaginitis is caused by several microbial agents and non-infective irritants. It is usually associated with urethritis, dysperunia, dysmenorrhea and lower abdominal pain. Microbial agents are transmitted through sexual contact and unhygienic habits. Hence, this study is designed to rule out the incidence of Microbial agents in reproductive women by microscopy, staining & culture. 100 patients after sensitization of the project were taken for this study Samples processed by microscopy, special staining and culture. The culture was performed with KUPFERBERG media. Out of 100 samples, 54 grew bacterial colonies, 30 were found to be positive for *Trichomonas vaginalis*, 2 for *Candida albicans* and the remaining 14 were having normal vaginal flora. The parasitic agents were found in 30 swabs by Giemsa stain. Preliminary wet mount microscopic examination revealed 15 cases to be positive. The analysis showed the correlation between bacterial and parasitic isolation, which was more common among sexually active female patients, which requires screening of patients in appropriate manner and treat accordingly to envisage a healthy quality life for the patients.

Introduction

Vaginitis is a known infectious disease caused by several microbial agents and non-infective irritants. The disease often involves vulva with mixed microbial agents. Therefore it is necessary to know the causative agents for patient care and management.

Vagina has mixed anaerobic and aerobic bacterial flora including *Gardnerella vaginalis* causing bacterial vaginosis. Infections with *Clostridium perfringens* typically after illegal abortions with non sterile instruments is also a well established entity (1).

Genital tract Infections of women are well known in reproductive age group and they are the commonest problem that makes a lady to visit her gynecologist with vaginal discharge. This can happen due to bacterial, fungal and parasitic microbial agents. It is usually associated with urethritis, dysperunia, dysmenorrhea and lower abdominal pain.

Trichomonas vaginalis is an important cause for sexually transmitted infection in reproductive age group. It is the most prevalent non- viral, non –bacterial cause for genital tract infection. It can also co-exist with HIV, Herpes simplex viruses, and can enhance the risk of Human Papilloma virus infection in women (2).

This disease is caused by a pear shaped protozoan *Trichomonas vaginalis*, giving rise to frothy, foul- smelling discharge with vulvo- vaginal irritation and lower abdominal pain (3).

It can be an important cause for Pelvic inflammatory Disease (PID) leading to complications like infertility (4).

In this era of infertility and sterility with long married life, we decided to take up this study among reproductive age group, among our patients attending OG clinic in our hospital.

The main aim of this study includes to know the incidence of Microbial agents in reproductive age group women attending our hospital with Leucorrhoea. Demonstration of the agent by microscopy, staining techniques. Demonstration of the agent by culture techniques. To demonstrate the sensitivity of culture isolation from clinical samples. And also to correlate the clinical findings.

Materials and Methods

100 willing patients after sensitization of the project, who were attending our OG-OPD were taken for this study.

Inclusion Criteria

Willing, Reproductive age group women with vaginal discharge.

Exclusion Criteria

Elderely (>50) and Young (<15), other vaginitis conditions.

Methods

A total of 100 samples were taken from 100 willing, sexually active female patients with an age group ranging from 20-50 years (Table: 1).

The vaginal swab with appropriate labelling and duly filled up request form was received in microbiology laboratory and subjected for microscopic examination, specific special staining and culture inoculation and incubation.

For Bacterial Etiology

The swab was inoculated onto the Mac Conkey agar, Blood agar and Chocolate agar and incubated at 37°C for 48 hours. The culture plates were examined for colony growth and processed up to species level identification and antibiotic sensitivity performed as per NCCLS guidelines (5).

For Parasitic STD

The direct wet mount of vaginal smear was examined by direct microscopic examination. The culture was performed

with KUPFERBERG media, though there are other media like, Feinberg, Whittington media. The swab was inoculated into the media incubated at 37°C for 7 days. The media was examined on 3rd, 5th, 7th day and a strict aseptic precautions. Both wet mount microscopic and Giemsa staining were done (Table: 2)

The culture media and Giemsa stains were performed as per NCCLS guidelines (6).

For fungal Causes

Samples were inoculated onto Sabourauds Dextrose Agar and fungal colonies grown were identified up to species level.

Results and Discussion

Out of 100 samples, 54 samples grew bacterial colonies, 30 samples were found to be positive for *Trichomonas vaginalis*, 2 samples were positive for *Candida albicans* and the remaining 14 were having normal vaginal flora.

Among the bacterial colonies 52 samples had Gram negative bacterial growth and 2 samples had Gram positive cocci.

The parasitic agents were found to be positive in 30 swabs and identified by culture and staining techniques by Giemsa stain. Preliminary wet mount microscopic examination revealed 15 cases to be positive (Table: 3).

2 samples yielded *Candida albicans* were found commonly in the 4th decade of age group.

Both bacterial, parasitic agents were found commonly in the 3rd decade of age group (Table: 4).

Gram positive bacteria, isolates were MSSE (Methicillin Sensitive *Staphylococcus epidermidis*). The gram negative isolates had an array of bacteria: *Escherichia coli*, *Klebsiella oxytoca* and *Klebsiella pneumonia*, *Pseudomonas aeruginosa* and *Acinetobacter baumannii*.

The KUPFERBERG culture media was more fruitful as it yielded positive results on 3rd day, 5th day, and 7th day proving to be more specific and sensitive for clinching the parasitic STD agent.

Totally they were 30 % culture positive for *Trichomonas vaginalis*, which was further confirmed by demonstration of the pear shaped protozoan by Giemsa staining technique (Pic:1)

The analysis of the 100 samples showed the following microbial profile (Table.5)

We got 54 bacterial vaginitis with heavy growth of *Escherichia coli*, *Klebsiella oxytoca*, *Klebsiella pneumonia*, *Pseudomonas aeruginosa*, *Staphylococcus epidermidis* and *Acinetobacter baumannii*. Their distribution is shown in (Table: 6)

The vaginal flora has complex microbiota with varying quantities and relative proportions (7).

There is an intricate balance among the microbes that maintains the normal vaginal flora. It is mainly dominated by the genus *Lactobacillus* that gives acidic pH to the vagina which guards against vaginitis.

When this pH gets altered, or colonization due to other bacteria and unhealthy sexual contact gives rise to vaginitis, which leads to vaginal discharge, urethritis, dysmenorrhea, dysperunia, lower abdominal pain and infertility.

Table.1 Age Distribution of the Patients (100)

AGE GROUP (YEARS)	PERCENTAGE %
20-25	20
26-30	21
31-35	30
36-40	15
41-45	07
46-50	04
51 & Above	04

Table.2 Culture Wet Mount Preparation and Giemsa Staining

S:NO	INCUBATION DAYS	MICROSCOPY CULTURE	STAIN
1	3 rd day	15	15
2	5 th day	10	10
3	7 th day	5	5

Table.3 Direct Wet Mount Preparation:

Diagnosis	NO. positive for <i>T. Vaginalis</i>	NO. negative for <i>T. Vaginalis</i>
Wet mount preparation	15	85

Table.4 Age Group Correlation with Parasitic STD


S.No	Age	Positive for <i>T.vaginalis</i>	Positive for Bacterial vaginitis
1	20-30	21	23
2	31-40	6	16
3	41-50	3	8
4	50 and above	-	7

Table.5 Microbial Profile of Vaginal Swab

S.No	Age	No.of Bacterial vaginitis positive	No.of <i>T. vaginalis</i> positive	No. of fungal vaginitis positive	No. of Normal vaginal flora
1	20-30	23	21	1	8
2	31-40	16	6	1	5
3	41-50	8	3	-	1
4	50 & above	7	-	-	-
		54	30	2	14

Table.6 Isolation Rates of Bacteria in Vaginal Swab

S.No	Name of the bacterial isolate	Sensitivity Pattern					
		Imi(S) %	Ak(S) %	G(S) %	Cz(S) %	Ca(S) %	Ce(S) %
1	<i>Escherichia coli</i> (19)	100	100	100	0	20	26
2	<i>Klebsiella oxytoca</i> (12)	100	100	100	0	20	26
3	<i>Klebsiella pneumonia</i> (10)	100	100	100	0	20	26
4	<i>Pseudomonas aeruginosa</i> (5)	100	100	94	0	15	22
5	<i>Acinetobacter baumannii</i> (2)	100	100	90	0	10	20
6	<i>Staphylococcus epidermidis</i> (2)	-	90	100	10	100	100



The analysis of the vaginal discharge with patients attending our OG department, gave rise to a wide range of microbial profile including bacterial, parasitic and fungal agents. About 54% of the samples had bacterial culture positivity, in which *Escherichia coli* was 39% positive (20/54) samples, *Klebsiella oxytoca* contributed for

24% with 13 samples, and *Klebsiella pneumoniae* had a share of 19% with (10/54) samples, the other isolates were *Pseudomonas aeruginosa* which was isolated from 5 samples contributing for 10% positivity and *Acinetobacter baumannii* and *Staphylococcus epidermidis* 4% with (2/54) samples in each. These bacterial

isolates were found in the vaginal swabs due to the lack of personal hygiene, as majority of our patients come from low socio economic status, with paddy field work pattern. which would have enhanced the colonization of this organism from the large intestine.

The next important isolate was parasitic agent *Trichomonas vaginalis* which contributed for 30%, was responsible for green coloured foul-smelling discharge, itching in vagina & vulva. It is the major sexually transmitted non-viral, non-bacterial disease, present all over the world.

It inhabits the genitalia, urinary tract in female, urethra and prostate in male. It takes shelter and nutrition from another living organism causing vulvo-vaginitis, PID. It can also precipitate infertility and sterility.

Kupferburg Media was selected as it contains selective agent to inhibit the growth of GNB & GPC. In addition it contains nitrogenous compounds for ensuring the growth of *Trichomonas vaginalis*. The inclusion of 0.1% agar reduces the oxygen tension for the prolific growth of *Trichomonas vaginalis*. Our study shows 30% of the samples were positive for *Trichomonas vaginalis* (Table: 2).

The sensitivity pattern of the gram negative bacilli showed 100% sensitivity to Imipenem, Amikacin and Gentamycin. 100% resistant for Cefazolin, 80% for Ceftazidime and 74% for Cefotaxime.

The analysis of Gram positive cocci (*Staphylococcus epidermidis*), had the following pattern, 100% oxacillin, linezolid and vancomycin sensitive. It had 50% resistant to Gentamycin and Ampicillin.

Other than the bacterial, parasitic agents,

the study also showed the presence of fungal agent in the sample- *Candida albicans* (2/100) with 2%. This also may be associated with sexual transmission similar to *Trichomonas vaginalis*.

In conclusion, the present study reveals that the vagina of the reproductive age group female gets inhabited with a variety of microorganisms due to faulty hygienic habits, which enhances migration of bacteria from intestine to genital tract.

Coitus is another precipitating factor for vaginitis which aids in the parasitic, fungal vaginitis. As a health professional it is mandatory to give health education to the patients with regards to personal hygienic habits and safe coitus in order to prevent vaginitis from happening in young reproductive ladies as shown in our studies.

This analysis shows the correlation between bacterial and parasitic isolation, which was more common among sexually active female patients as shown in our study, which requires screening of patients in appropriate manner and treat accordingly to envisage a healthy quality life for the patients.

References

1. Koneman's. Colour Atlas and Textbook of Diagnostic Microbiology, 6th Edition, 88-90 P.
2. Lucia Pattullo, Sarah Griffeth, Lili Ding et al. Stepwise Diagnosis of *Trichomonas vaginalis* Infection in Adolescent Women. J. Clin. Microbiol 2009; 47: 59-63.
3. E. D. Jatau, O. S. Olonitola and A. T. Olayinka. Prevalence of *Trichomonas* Infection among Women Attending Antenatal Clinics in Zaria, Nigeria. Annals of African Medicine 2006; 5: 178-181.

4. Nashaat E. Nassel, Amara F. Afif et al. Evaluation of microscopy and polymerase chain reaction for diagnosis of symptomatic and asymptomatic female Trichomoniasis. *Egyptian Parasitologists United Society Journal* 2014; 7: 37-46.
5. National committee for clinical Laboratory Standards. Abbreviated identification of Bacteria and Yeast; Approved Guidelines. NCCLS document M35-A. Wayne, PA: NCCLS, 2002.
6. Z. T. Valadkhani, M. Assmar, N. Hassan et al. The Prevalence of Trichomoniasis in High-Risk Behavior Women Attending the Clinics of Tehran Province Penitentiaries. *Iran J Med Sci* 2010; 35(3) :190-194.
7. Shamim Mumtaz, Mumtaz Azhmed et al. Aerobic vaginal Pathogens and Their Sensitivity Pattern. *J Ayub Med Coll Abbottabad* 2008; 20(1): 113-117.

How to cite this article:

Karthika Jayakumar, Wills shiela, S. Jayalakshmi, Sathya Pandurangan and M. Mohanambal. 2016. Microbial Vaginitis in Reproductive Women of Silk city of South India. *Int.J.Curr.Microbiol.App.Sci*.5(2): 272-278. doi: <http://dx.doi.org/10.20546/ijcmas.2016.502.031>

Majority of the women present to the gynecology OPD with complains of Senior Resident, Department of white discharge, on failure of recognizing the yeast at the appropriate time, leads to frequent hospital visit Obstetrics and Gynecology, ESIC and lower genital discomfort. Reproductive tract importance of association of pregnancy, preterm labor and intra infections among young married women in Tamil Nadu, uterine device with candidiasis strongly indicates the need of sex India, International Family Planning Perspectives. A study of high vaginal swabs in tract infection isolated in Benin City, Edo state, Nigeria Candida albicans and non albicans species as 6. Van Dyck E, Meheus AS, Piot P. A manual on Laboratory etiological agent of vaginitis in pregnant women Vaginitis amongst Women of Reproductive Age Group from. Navi Mumbai, India. Anuradha Narayankhedkar, 1Anahita Hodiwala, 2and Arati Mane3. It is noteworthy that the mixed microbial etiologies observed. were all seen in association with BV. It may thus be speculated. vaginosis in women with vaginal symptoms in South Province, Rwanda, African Journal of Clinical and Experimental Microbiology, vol. 1, no. 1, pp. 1-4. Candida species are one of the commonest causes of vaginitis in healthy women of reproductive age. Vulvovaginal candidiasis (VVC) is characterized by vulvovaginal itching, redness and discharge.